

YIELD AND CHEMICAL COMPOSITION OF BREAD WHEAT CULTIVARS AS AFFECTED BY SOME SKIPPING IRRIGATION

S. A. S. Mehasen⁽¹⁾, N. Kh. El-Gizawy⁽¹⁾, A. M. Sharoba⁽²⁾, S. A. Soliman⁽³⁾
and T. R. M. Khalil⁽⁴⁾

⁽¹⁾ Agron. Dep., Fac. of Agric., Moshtohor, Benha Univ., Egypt.

⁽²⁾ Food Sci. Dep., Fac. of Agric., Moshtohor, Benha Univ., Egypt.

⁽³⁾ Institute of Food technology, Agricultural Research Center, Giza, Egypt.

⁽⁴⁾ Food specialist, Ain Shams Univ., Egypt.

(Received: Apr. 23 , 2014)

ABSTRACT: *Two field experiments were conducted at the Experiment and Research Center, Fac. Agric., Moshtohor, Benha Univ., Kalubia Governorate, Egypt, during the two growing seasons 2010/11 and 2011/12 seasons, to study the effect of five skipping an irrigation at tillering stage, elongation stage, heading stage, milk stage and filling stage on yield and chemical properties of five bread wheat cultivars (Giza 168, Sakha 94, Gemmaza 9, Sids 13 and Misr 1). The obtained results were as follows:*

Skipping an irrigation at tillering, elongation and heading stages decreased flag leaf area, plant height, number of spikes m^{-2} , 1000-kernel weight, grain yield and straw yield compared with skipping irrigation at filling stage treatments. Moreover, skipping irrigation at tillering stage was earlier heading compared with other skipping irrigation treatments.

Wheat cultivars were significantly different in all traits studied. Gemmaza 9 cultivar gave the highest values of flag leaf area, plant height, spike length, No. of spikelets spike⁻¹, No. of spikes m^{-2} , seed index, grain yield and straw yield, also Gemmaza 9 cultivar was earlier heading (89.67 day) compared with other wheat cultivars.

Significant effect of interaction between skipping irrigation and wheat cultivars was obtained for all growth, yield and yield components except seed index. Skipping irrigation at tillering stage for Gemmaza 9 cultivar gave the highest value of No. of spikelets spike⁻¹ and surpassed the other 4 cultivars in early heading. While, skipping irrigation at milk stage for Gemmaza 9 cultivar gave the highest values of spike length and No. of spikes m^{-2} . Moreover, skipping irrigation at filling stage for Gemmaza 9 cultivar gave the highest values of flag leaf area, plant height, grain yield fed^{-1} and straw yield fed^{-1} .

Skipping an irrigation at milk and filling stages decreased moisture, fat and carbohydrate contents, whereas, the same two treatments gave the highest values of protein, ash and fiber contents in wheat kernels compared with other irrigation treatments. The lowest values of moisture and carbohydrate contents were obtained by Giza 168 cultivar and also, the same cultivar gave the highest value of protein content, while, Sids 13 cultivar gave the highest value of ash content, Sakha 94 cultivar gave the highest value of fiber content and Misr 1 cultivar gave the highest value of fat content and in wheat kernels.

Skipping irrigation at filling stage for Giza 168 cultivar gave the lowest values of moisture and carbohydrate contents, also, the same treatment gave the highest value of protein content in wheat kernels. While, skipping irrigation at milk stage for Sids 13 cultivar gave the highest value of ash content and skipping irrigation at filling stage for Sids 13 cultivar gave the highest value of fiber contents. Whereas, skipping irrigation at milk or tillering stages for Giza 168 or Gemmaza 9 cultivars gave the highest value of fat content.

Keywords: *Wheat, Cultivars, Skipping irrigation, Yield and Chemical composition.*

INTRODUCTION

Wheat is considered the main source of food in the world and in Egypt. Raising wheat production through increasing

productivity and increasing the cultivated area is an important national target to minimize the gap between the Egyptian production and consumption. In Egypt, the

Yield and chemical composition of bread wheat cultivars as affected.....

wheat cultivated area is about 3.16 million feddans producing 8.795 million tons (Agric. Statistic Bulletin, 2012). Increasing wheat yield per unit area can be achieved by breeding high yielding varieties or improving the cultural treatments of the crop. Modern wheat varieties were developed to maximize grain yield under favorable environmental conditions (high input conditions especially water supply). In the light of the present national water policy concerning saving irrigation water expanding wheat area needs more searching for varieties produce high yield under suitable water regime. But safe saving an irrigation, beside it closely dependence on plant growth stage, is greatly influencing by a number of factors especially cultivars.

Many investigators had reported the effect of skipping an irrigation at deferent growth stages of wheat on yield and its components as well as chemical properties (El-Kalla *et al*, 1992; Abd El-Gawad *et al*, 1993; El-Monoufi and Harb, 1994; Abo-Shetaia and Abd El-Gawad, 1995; Ghandorah *et al*, 1997; Bayoumi, 1999; Eman, 2000; Sharaan *et al*, 2000; Abd-AllaKotb, 2005 Ramadan and Awaad 2008 and El-Nady, 2009).

Several investigators showed that wheat cultivars differed in yield and its components as well as chemical properties (Mehasen, 1999; El-Hawary, 2000; Sharaan *et al*, 2000; Toaima *et al*, 2000; Abd El-hameed, 2002; Ali *et al*, 2004; Mehasen and Mohamed, 2005; Abu-Grab *et al*, 2006; Omar, 2007; El-Ganayniand and Mahmoud, 2008; Hassan, 2008; El-Nady, 2009; Mehasen *et al*, 2009; Ashmawy *et al*, 2010; Abd El-Nour and Fateh, 2011; Mehasen *et al*, 2013 and Badwee, 2014).

Sharaan *et al*, (2000) studied the performance and productivity of five wheat varieties (i.e., Sids 1, Sakha 8, Sakha 69, Giza 164, Giza 167) under three water regimes (i.e., skipping one irrigation at heading stage, skipping one at dough stage, and normal six irrigations). Significant interaction effects between water regimes and wheat varieties on plant height, number and weight of grains/spike and 1000-grain

weight were detected in the combined analysis.

Therefore, the present investigation was designed to study the performance and productivity of bread wheat cultivars under different skipping an irrigation at growth stages, in Kalubia Governorate.

MATERIALS AND METHODS

Two field experiments were conducted at the Experimental Research Center, Faculty of Agriculture at Moshtohor, Benha Univ., Kalubia Governorate, Egypt, during the two successive growing seasons 2010/11 and 2011/12, to study the effect of five skipping an irrigation attillering stage, elongation stage, heading stage, milk stage and filling stage on yield and chemical properties of five bread wheat cultivars (Giza 168, Sakha 94, Gemmaza 9, Sids 13 and Misr 1). The soil was clay in texture with a pH value of 7.80, 7.78 and an organic matter content of 1.66, 1.75% and available N of 52, 55 ppm during the two growing seasons, respectively.

The treatments were assigned in a split-plot design with four replications. Irrigation treatments were arranged at random in the main plots while, cultivars of wheat occupied the sub-plots. The sub-plot area was 10.5 m² consisted of 15 rows of 3.5 m length and spaced 20 cm apart. Wheat cultivars were cultivated on November 25th and 27th in the first and second seasons, respectively. The preceding crop was maize in both seasons. The normal cultural practices were carried out as recommended in the area.

Studied characters:

The studied traits were No. of days to 50% heading and flag leaf area (cm²). Random samples of 10 guarded plants were taken from sub-plots at harvesting time to determine the following characters: plant height (cm), spike length (cm), and No. of spikelet spike⁻¹. For determining No. of spikes/m² and seed index (g) a sample of one square meter from each sub-plot was taken. Grain and straw yields (kg fed⁻¹) were estimated on whole sub-plot basis.

Chemical analysis:

Moisture, protein, ash, fiber, fat and carbohydrates were determined according to the methods recommended by A.O.A.C. (2000).

Statistical analysis:

Analysis of variance was done for the data of each season separately and the combined analysis of variance for two seasons was conducted testing the error homogeneity according to Snedecor and Cochran (1980) treatment means were compared using least significant difference test at 0.05 level of significance. Using the MSTAT-C Statistical Software package (Michigan State University, 1983)

RESULTS AND DISCUSSION

-Effect of skipping irrigations.

The mean values of the traits studied as affected by irrigation treatments are presented in Table (1). Skipping an irrigation at tillering, elongation, heading, milk and filling stages treatments showed significant differences in all traits studied in the combined analysis. Skipping one irrigation either at tillering or at elongation or at heading stages decreased all traits studied except No. of days to 50% heading, spike length, No. of spikelets spike⁻¹ and seed index. Skipping an irrigation at tillering, elongation and heading stages decreased flag leaf area, plant height, number of spikes m⁻², 1000-kernel weight, grain yield and straw yield by 15.54, 9.85 and 2.97%, 3.37, 2.58 and 2.01%, 54.16, 50.02 and 11.26%, 5.58, 4.38 and 4.66%, 13.00, 10.38 and 7.10% and 21.43, 18.74 and 14.86% respectively, compared with skipping irrigation at filling stage treatments. Moreover, skipping irrigation at tillering stage was earlier heading compared with other skipping irrigation treatments. The negative effect of yield and its components caused by skipping an irrigation could be explained on the basis of the loss of turgor which affects the rate of cell expansion and ultimate cell size. Loss of turgor is probably the most sensitive process to water stress, thus, decrement in growth rate, stem elongation and leaf expansion. In this

connection, El-Kalla *et al*, (1992); Abd El-Gawad *et al*, (1993); El-Monoufi and Harb (1994); Abo-Shetaia and Abd El-Gawad (1995); Ghandorah *et al*, (1997); Bayoumi (1999); Eman (2000); Sharaan *et al*, (2000); Abd-AllaKotb, (2005) and El-Nady (2009), reached the same conclusion.

-Cultivars differences.

The results reported in Table (1) indicate clearly that, there were significant differences between the different wheat cultivars in all traits studied. Moreover; it is clear from Table (1) that Gemmaza9 cultivar gave the highest values of flag leaf area (48.52 cm²), plant height (106.57 cm), spike length (14.00 cm), No. of spikelets spike⁻¹ (20.02 spikelets), No. of spikes m⁻² (335.6 spike), seed index (44.45 gm), grain yield (3376.8 kg fed⁻¹) and straw yield (3986.7 kg fed⁻¹). Also Gemmaza9 cultivar was earlier heading (89.67 day) compared with other wheat cultivars. On the other hand, Giza 168 cultivar gave the lowest values of flag leaf area (44.90 cm²), plant height (89.82 cm), spike length (10.82 cm), No. of spikelets spike⁻¹ (19.00 spikelets), No. of spikes m⁻² (300.0 spike), seed index (39.11 gm), grain yield (2442.3 kg fed⁻¹) and straw yield (3385.2 kg fed⁻¹). Also Giza 168 was latest heading (92.45 day) compared with other wheat cultivars. It could be concluded that varietal differences among wheat cultivars may be due to genetical make up. The superiority of Gemmaza 9 cultivar in grain yield (kg fed⁻¹) over other wheat cultivars might be due to the increase in yield components, namely, spike length, No. of spikelets spike⁻¹ and seed index. The results obtained by Mehasen (1999); El-Hawary (2000); Toaima *et al*, (2000); Abd El-Hameed (2002); Ali *et al*, (2004); Mehasen and Mohamed (2005); Abu-Grab *et al*, (2006); Omar (2007); El-Ganayni and Mahmoud (2008); Hassan (2008); Mehasen *et al*, (2009); Ashmawy *et al*, (2010); Abd El-Nour and Fateh (2011); Mehasen *et al*, (2013) and Badwee (2014) indicated marked differences among wheat varieties in growth, yield and yield components.

Yield and chemical composition of bread wheat cultivars as affected.....

Table 1. Growth, yield and yield components of wheat as affected by skipping irrigation and Cultivars (Combined analysis of 2010/11 and 2011/12 seasons)

Treatments	No. days to 50% heading	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	No. spikelet spike ⁻¹	No. spikes m ⁻²	Seed index (g)	Grain yield (kg fed ⁻¹)	Straw yield (kg fed ⁻¹)
Skipping at									
Tillering stage	89.88	43.11	96.31	12.74	19.50	246.1	41.19	2625.9	3264.0
Elongation stage	90.39	44.90	98.02	12.98	19.42	252.9	41.30	2688.5	3338.0
Heading stage	91.02	46.83	99.57	11.83	18.95	341.0	40.83	2770.7	3450.8
Milk stage	91.65	48.50	101.02	12.74	19.47	355.1	42.03	2867.4	3983.3
Filling stage	92.21	49.18	102.18	12.74	19.67	379.4	43.11	2967.5	4013.7
L.S.D at 5%	0.20	0.19	0.51	0.08	0.31	3.37	0.51	24.3	87.9
Cultivars									
Giza 168	92.45	44.90	89.82	10.82	19.00	300.0	39.11	2442.3	3385.2
Sakha 94	90.66	47.66	99.15	13.05	19.55	333.5	42.20	2931.2	3704.7
Gemmaza 9	89.67	48.52	106.57	14.00	20.02	335.6	44.45	3376.8	3986.7
Sids 13	91.60	45.61	100.90	13.20	19.30	300.8	41.25	2560.2	3648.7
Misr 1	90.96	46.46	99.00	11.96	19.15	300.6	41.46	2609.6	3274.3
L.S.D at 5%	0.16	0.19	0.68	0.10	0.25	3.20	0.35	17.2	96.9

-Interaction effect:

Significant effect of interaction between skipping irrigation and wheat cultivars was obtained for all growth, yield and yield components except seed index namely, No. of days to 50% heading, flag leaf area, plant height, spike length, No. of spikelet spike⁻¹, No. of spikes m⁻², grain and straw yields fed⁻¹ (Table 2). Skipping irrigation at tillering stage for Gemmaza 9 cultivar gave the highest value of No. of spikelets spike⁻¹ (10.87 spikelet) and surpassed the other 4 cultivars in early heading (88.02 day). While, skipping irrigation at milk stage for Gemmaza 9 cultivar gave the highest values of spike length (14.36 cm) and No. of spikes m⁻² (398.9 spikes). Moreover, skipping irrigation at filling stage for Gemmaza 9 cultivar gave the highest values of flag leaf area (51.45 cm²), plant height (109.75 cm), grain yield fed⁻¹ (3591.6 kg) and straw yield fed⁻¹ (4473.3 kg). On the other hand, skipping irrigation at tillering stage for Giza

168 cultivar gave the lowest values of flag leaf area (41.45 cm²), plant height (84.37 cm), No. of spikes m⁻² (221.8 spike), grain yield fed⁻¹ (2289.3 kg) and straw yield fed⁻¹ (2975.6 kg). Skipping irrigation at heading stage for Giza 168 cultivar gave the lowest values of spike length (10.53 cm) and No. of spikelets spike⁻¹ (18.12 spikelet) while, skipping irrigation at filling stage for Giza 168 cultivar gave the latest heading (94.01 day). Similar results were also reported by Sharaan *et al.*, (2000) and El-Nady (2009).

-Chemical properties of wheat kernels:

Twenty five of wheat kernels samples (five skipping an irrigation at tillering, elongation, heading, milk and filling stages treatments and five bread wheat cultivar i.e. Giza 168, Sakha 94, Gemmaza 9, Sids 13 and Misr 1), were analyzed for their chemical composition, i.e., moisture, ash, protein, fat,

Yield and chemical composition of bread wheat cultivars as affected.....

crude fiber and carbohydrate contents. The obtained results are shown in Tables (3&4).

Table 2. Effect of the interaction between Skipping irrigation and Cultivars on growth, yield and yield components of wheat (over the combined analysis)

Treatments		No. days to 50% heading	Flag leaf area (cm ²)	Plant height (cm)	Spike length (cm)	No. spikelets Spike ⁻¹	No. spikes m ⁻²	Grain yield (kg fed ⁻¹)	Straw yield (kg fed ⁻¹)
Skipping at	Cultivars								
Tillering stage	Giz 168	91.07	41.45	84.37	11.02	19.00	221.8	2289.3	2975.6
	Sakh 94	89.66	43.99	98.50	13.07	19.25	275.6	2748.0	3374.5
	Gem 9	88.02	44.85	102.37	14.18	20.87	270.9	3198.3	3734.1
	Sids 13	90.85	42.42	98.37	13.35	19.37	236.3	2454.8	3260.1
	Misr 1	89.82	42.83	97.75	12.08	19.00	225.8	2439.0	2976.0
Elongation stage	Giz 168	92.03	43.14	88.87	10.76	19.12	223.3	2351.2	3243.7
	Sakh 94	89.89	45.97	98.00	13.45	19.75	276.1	2829.3	3140.6
	Gem 9	88.75	46.63	102.75	14.62	19.62	278.4	3285.2	3787.6
	Sids 13	91.04	43.85	102.37	13.80	19.25	244.4	2452.5	3392.5
	Misr 1	90.22	44.92	98.12	12.26	19.37	242.5	2524.3	3125.6
Heading stage	Giz 168	92.05	45.02	94.37	10.53	18.12	338.8	2424.3	3025.6
	Sakh 94	90.00	48.02	97.12	12.13	19.12	363.9	2918.7	3721.2
	Gem 9	90.03	48.46	109.25	13.26	19.50	364.5	3356.2	3631.2
	Sids 13	92.98	46.27	99.50	12.20	18.87	322.3	2537.5	3332.5
	Misr 1	90.04	46.40	97.62	11.02	19.12	315.8	2616.6	3293.3
Milk stage	Giz 168	93.12	46.98	90.12	10.87	19.37	347.1	2521.1	4190.5
	Sakh 94	90.66	49.20	98.87	13.15	19.37	375.9	3066.2	4186.2
	Gem 9	90.42	51.22	108.75	14.36	20.50	398.9	3452.5	4207.5
	Sids 13	92.10	47.15	104.12	13.10	19.12	320.0	2624.3	3965.6
	Misr 1	91.95	47.97	103.25	12.21	19.00	333.8	2673.1	3416.8
Filling stage	Giz 168	94.01	47.93	91.37	10.90	19.37	369.3	2625.3	3490.8
	Sakh 94	93.10	51.15	103.25	13.47	20.25	376.0	3093.7	4221.2
	Gem 9	91.15	51.45	109.75	13.56	19.62	385.5	3591.6	4473.3
	Sids 13	91.05	48.34	100.12	13.56	19.87	381.1	2731.8	4273.1
	Misr 1	91.76	50.20	98.25	12.23	19.25	385.0	2795.0	3560.0
L.S.D at 5%		0.35	0.43	1.52	0.21	0.56	7.01	38.5	216.7

Table 3. Chemical composition of wheat kernels as affected by skipping irrigation and Cultivars (Combined analysis of 2010/11 and 2011/12 seasons)

Treatments	Moisture%	As a dry weight basis%				
		Protein	Ash	Fiber	Fat	Carbohydrates
Skipping at						
Tillering stage	9.68	11.33	1.64	2.37	2.06	82.60
Elongation stage	9.60	11.14	1.56	2.47	2.10	82.73
Heading stage	9.28	11.26	1.52	2.42	2.14	82.66
Milk stage	8.56	12.24	1.94	3.00	1.50	81.32
Filling stage	8.58	12.11	1.56	3.13	1.18	82.02
Cultivars						
Giza 168	8.75	12.99	1.56	2.64	1.82	80.99
Sakha 94	9.56	11.76	1.62	2.90	1.92	81.80
Gemmaza 9	9.24	11.92	1.54	2.63	1.90	82.01
Sids 13	8.88	10.56	1.82	2.74	1.42	83.46
Misr 1	9.24	10.87	1.78	2.47	2.00	82.88

Yield and chemical composition of bread wheat cultivars as affected.....

Table 4. Chemical composition of wheat kernels as affected by skipping irrigation and cultivars (Combined analysis of 2010/11 and 2011/12 seasons)

Treatments		Moisture %	As a dry weight basis%				
Skipping at	Cultivars		Protein	Ash	Fiber	Fat	Carbohydrates
Tillering stage	Giza 168	9.0	12.80	1.6	2.20	2.5	80.90
	Sakha 94	9.9	11.49	1.6	2.73	2.2	80.98
	Gemmaza 9	9.5	11.97	1.5	2.23	2.2	82.10
	Sids 13	10.0	10.00	1.8	2.25	1.6	84.35
	Misr 1	10.0	10.39	1.7	2.46	1.8	83.65
Elongation stage	Giza 168	9.8	11.87	1.4	2.50	2.2	82.03
	Sakha 94	9.7	11.32	1.5	2.90	2.1	82.18
	Gemmaza 9	9.8	11.72	1.5	2.24	2.5	82.04
	Sids 13	8.7	10.15	1.7	2.31	1.5	84.34
	Misr 1	10.0	10.64	1.7	2.38	2.2	83.08
Heading stage	Giza 168	8.5	13.00	1.5	2.40	1.9	81.20
	Sakha 94	9.8	11.21	1.6	2.70	2.5	81.99
	Gemmaza 9	9.4	11.51	1.4	2.38	1.9	82.81
	Sids 13	9.5	10.13	1.6	2.31	1.9	84.06
	Misr 1	9.2	10.43	1.5	2.31	2.5	83.26
Milk stage	Giza 168	8.6	13.41	1.8	3.00	1.2	80.59
	Sakha 94	9.2	12.35	2.0	2.90	1.6	80.60
	Gemmaza 9	8.7	12.54	1.7	3.00	1.6	81.16
	Sids 13	7.8	11.20	2.3	3.38	1.2	82.01
	Misr 1	8.5	11.36	1.9	2.61	1.9	82.23
Filling stage	Giza 168	8.0	13.91	1.5	3.10	1.3	80.19
	Sakha 94	9.2	11.43	1.4	3.20	1.2	82.77
	Gemmaza 9	8.8	11.87	1.6	3.30	1.3	81.93
	Sids 13	8.4	11.33	1.7	3.46	0.9	82.61
	Misr 1	8.5	11.51	1.6	2.61	1.6	82.68

Data in Table (3) cleared that the lowest values of moisture contents and carbohydrate contents in wheat kernels were 8.60, 8.54% and 81.32, 82.02% were obtained for skipping irrigation at milk stage and skipping irrigation at filling stage treatments, respectively. Also, the same two treatments gave the highest value of protein content in wheat kernels were 12.24 and 12.11% were obtained for skipping irrigation at milk stage and skipping irrigation at filling stage treatments, respectively. Moreover, skipping irrigation at milk stage treatment gave the highest value of ash content in wheat kernels was 1.94% compared with other skipping irrigation treatments. Whereas, skipping irrigation at filling stage treatment gave the highest value of fiber content in wheat kernels was 3.13% compared with other skipping irrigation treatments. On the other hand, skipping an irrigation at heading, elongation, and tillering

stages treatments gave the highest value of fat content in wheat kernels were 2.14, 2.10 and 2.06% respectively, compared with other skipping irrigation treatments. Moisture, fat and carbohydrate contents decreased when irrigation deficiency treatments were done during the last two periods of plant age (milk and filling stages) while, protein, ash and fiber contents increased when irrigation deficiency treatments were done during the last two periods of plant age (fruiting and maturing stage).

Data also indicated that the lowest values of moisture contents and carbohydrate contents in wheat kernels were 8.71 and 80.99% respectively, were obtained by Giza 168 cultivar compared with other cultivars. Also, the same cultivar gave the highest value of protein content in wheat kernels (12.99%) compared with other cultivars.

Yield and chemical composition of bread wheat cultivars as affected.....

While, Sids 13 cultivar gave the highest value of ash content(1.82%), Sakha 94 cultivar gave the highest value of fiber content(2.90%) and Misr 1 cultivar gave the highest value of fat content in wheat kernels (2.00%) compared with other cultivars.

From the results presented in Table (4), it could be noted that moisture contents of wheat samples ranged from 7.8 to 10.0%, protein from 10.0 to 13.91 %, ash from 1.4 to 2.3%, fiber from 2.20 to 3.46%, fat from 0.9 to 2.5% and total Carbohydrates from 80.19 – 84.35%. Skipping irrigation at filling stage for Giza 168 cultivar gave the lowest values of moisture and carbohydrate contents in wheat kernels were 7.8 and 80.19% respectively. Also, the same treatment gave the highest value of protein content in wheat kernels (13.91%). While, skipping irrigation at milk stage for Sids 13 cultivar gave the highest value of ash contents (2.3%) and skipping irrigation at filling stage for Sids 13 cultivar gave the highest value of fiber contents (3.46%). Whereas, skipping irrigation at milk or tillering stages for Giza 168 or Gemmaza 9 cultivars gave the highest value of fat contents (2.5%).

REFERENCES

- AbdAllaKotb, M. (2005). Effect of foliar application of glycinebetaine on growth and yield of wheat (*Triticumaestivium* L.) under water stress. The 11th conference of Agronomy, Agron., Dept., Fac. Agric., Assiut Univ., Nov. 15-16.
- Abd El-Gawad, A. A., S. EL-Habbal, A. S. A. Edris and Elham A. Dorgham (1993). Effect of water stress during filling period and nitrogen fertilization on yield and its attributes of two wheat genotypes. Egypt J. Agron., 18(1-2):211-227.
- Abd El-Hameed, I. A. (2002). Effect of some agronomic practices on wheat. Ph. D. Thesis, Fac. Agric., Zagazig Univ., Egypt.
- Abd El-Nour, Nadya, A. R. and Hayam S. A. Fateh (2011). Influence of sowing date and nitrogen fertilization on yield and its components in some bread wheat genotypes. Egypt. J. Agric. Res, 89(4):1413-1433.
- Abo Shetaia, A. M. and A. A. Abd EL-Gawad (1995). Growth, yield and yield attributes of wheat in relation to N fertilization and withholding an irrigation at different stages of growth. Annals Agric. Sci., Fac. of Agric., Ain Shams Univ., Cairo, Egypt, 40 (1): 195-211.
- Abu-Grab, O. S., A. M. Mousa and G. A. El-Shaarawy (2006). Photosynthetic and N-use efficiencies for some wheat cultivars in relation to planting density and nitrogen fertilization level. Egypt. J. Appl. Sci., 21(2B): 475-492.
- Ali, G. A., O. E. Zeiton, A. H. Bassiouny and A. R. Y. A. El-Banna (2004). Productivity of wheat cultivars grown at El-Khattara and El-Arish under different levels of planting densities and N-fertilization. Zagazig J. Agric. Res., 31(4A): 1225-1256.
- A.O.A.C. (2000). Official Methods of Analysis Association of official analysis chemists, 14th Ed., Washington, D.C., USA.
- Ashmawy, F., M. S. El-Habal, H. S. Saoudy and ImanKh. Abbas (2010). The relative contribution of yield components to grain yield of some wheat cultivars grown under different nitrogen fertilizer levels. Egypt. J. Agric. Res., 88(1):225-239.
- Badwee, H. M. S. M. (2014). Analysis of response curve of wheat yield to nitrogen fertilization as related to environment pollution. Ph. D. Thesis, of Environ. Sci., Dept. of Agric. Sci., Inst. of Enviro., Ain Shams Univ., Egypt.
- Bayoumi, Y. T. (1999). Variability in morphophysiological characters and its relation to drought tolerance in some field crops. Ph. D. Thesis, Agron. Dept., Fac. Agric., Suez Canal Univ., Egypt.
- El-Ganayni, A. A. and Gamalat A. Mahmoud (2008). Productivity of two bread wheat cultivars under three seeding rates. J. Agric. Sci., Mansoura Univ., 33(6): 3993-4000.
- El-Hawary, M. A. (2000). Evaluation of some wheat varieties under water deficit conditions. Zagazig J. Agric. Res., 27(4): 819-830.
- EL-Kalla, S. E., A. A. Kandil, M. S. Sultan, S. A. EL-Moursy and A. M. Abd EL-Aal (1992). Response of some wheat

- varieties to irrigation regimes and time of nitrogen fertilization. *J. Agric. Sci. Mansoura Univ.* 17(8): 2568-2575.
- EL-Monoufi, M. M. and O. M. S. Harb (1994). Grain yield and yield components of wheat as effected by water stress at different plant stages. *Zagazig J. Agric. Res.*, 21 (4): 1023-1028.
- El-Nady, Hanim E. A. (2009). Breeding bread wheat of drought and earliness. Ph. D. Thesis, Fac. Agric., Moshtohor, Benha Univ., Egypt.
- Eman, S. S. (2000). Yield and yield attributes of wheat in relation to nitrogen and withholding an irrigation at different stages of growth. *Annals Agric. Sci., Fac. of Agric., Ain Shams Univ., Cairo*, 45(2): 439-452.
- Ghandorah, M. O., I. I El-Shawaf, Kh. A. Moustafa and A. M. Gadalla (1997). Evaluation of some early generation of bread wheat genotypes grown under heat and water stress at the central region of Saudi Arabia. *Arab. Gulf. J. Sci. Res.*, 15(2): 505-523.
- Hassan, Manal A. (2008). Effect of seeding rate and row spacing on productivity and resistance to powdery mildew of two bread wheat cultivars. *Egypt. J. Appl. Sci.*, 23(10A): 169-182.
- Mehasen, S. A. S and N. A. Mohamed (2005). Multivariate and response curve analyses for yield and its attributes in some wheat cultivars under nitrogen fertilization levels. *Egypt. J. Appl. Sci.*, 20(2): 93-109.
- Mehasen, S. A. S. (1999). Response of some wheat varieties to agrispn foliar application and nitrogen rates. *Annals of Agric. Sci., Moshtohor*, 37(2): 853 – 864.
- Mehasen, S. A. S., M. A. Ahmed and M. A. M. Morsy (2009). Evaluation of some wheat genotypes under different seeding rates. *Annals of Agric. Sci., Moshtohor*, 47 (3):167-174.
- Mehasen, S. A. S., M. A. Ahmed and S. Sh. Abdullah (2013). Improving productivity of bread wheat genotypes by using of some growth promoters. *Egypt. J. Plant Breed.*, 17(2):4-15.
- Michigan State University (1983). MSTAT-C: Micro- computer Statistical Program, Version 2.0. Michigan State University, East Lansing.
- Omar, A. E. A. (2007). Productivity of some wheat cultivars as affected by sowing date and seeding rate. *Egypt. J. Appl. Sci.*, 22(3): 103-116.
- Ramadan, A. R. and S. S. Awaad (2008). Response of yield attributes of some bread wheat varieties to irrigation levels and seeding rates under old land conditions. *J. Agric. Sci., Mansoura Univ.*, 33(7): 4717-4737.
- Sharaan, A.N., F.S. Abd EL-Samie and I.A. Abd EL-Gawad (2000). Response of wheat varieties to some environmental influences. II-Effect of planting date and drought at different plant stages on yield and its components. 9th Conf. of Agronomy 2-3 September, Menofiya, Egypt : 1-16.
- Snedecor, G. W. and W. G. Cochran (1980). *Statistical Methods*, 7th Ed., Iowa State Univ. Press, Ames, Iowa, USA.
- Toaima, S.E., A.A. El-Hofi and H. Ashoub (2000). Yield and technological characteristics of some wheat varieties as affected by nitrogen fertilizer and seed rates. *J. Agric. Sci., Mansoura Univ.*, 25(5): 2449-2467.

تأثير حرمان بعض أصناف قمح الخبزمن بعض الريات علي المحصول والصفات الكيميائية

صديق عبد العزيز صديق محيسن⁽¹⁾، ناصر خميس الجيزاوي⁽¹⁾، أشرف مهدي شروبة⁽²⁾،
سيد عبد الحميد سليمان⁽³⁾، طه رمضان مهدي خليل⁽⁴⁾

Yield and chemical composition of bread wheat cultivars as affected.....

(¹) قسم المحاصيل- كلية الزراعة بمشتهر - جامعة بنها- مصر .

(²) قسم علوم الاغذية - كلية الزراعة بمشتهر - جامعة بنها- مصر .

(³) معهد تكنولوجيا الأغذية - مركز البحوث الزراعية - الجيزة - مصر .

(⁴) أخصائي تغذية بجامعة عين شمس - مصر .

المخلص العربي

أقيمت تجربتان حقليتان بمركز البحوث والتجارب بكلية الزراعة بمشتهر - جامعة بنها- محافظة القليوبية- مصر خلال موسمي الزراعة ٢٠١٠/١١ و ٢٠١١/١٢ م لدراسة تأثير خمس معاملات لتحريم رية في مراحل مختلفة من حياة النبات وهي (تحريم رية في طور التفريع، تحريم رية في طور الاستطالة، تحريم رية في طور طرد السنابل، تحريم رية في طور البني للحبوب و تحريم رية في طور العجيني للحبوب) علي المحصول والصفات الكيميائية لخمسة أصناف من القمح الخبز وهي (جيزة ١٦٨، سخا ٩٤، جميزة ٩ ، سدس ١٣ ومصر ١) -ويمكن تلخيص أهم نتائج التحليل المشترك للموسمين فيما يلي.

- أدت معاملات تحريم رية في أطوار التفريع والاستطالة و طرد السنابل الي نقص معنوي في صفات مساحة ورق العلم (سم^٢) ، طول النبات (سم) ، عدد السنابل/م^٢ ، وزن ال ١٠٠٠ حبة (جم) ، محصول الحبوب ومحصول القش (كجم/فدان). وكذلك أدت معاملة تحريم رية في طور التفريع الي التباين في صفة طرد ٥٠% من السنابل. كما ادت معاملات تحريم رية في طوري الاستطالة و طرد السنابل الي نقص معنوي في صفتي طول السنبله (سم) وعدد ابراج السنبله عن بقية معاملات الري الاخرى.

- كانت هناك اختلافات معنوية بين الاصناف في جميع الصفات المدروسة. سجل صنف جميزة ٩ أعلى متوسطات لكل من مساحة ورقة العلم (سم^٢) ، طول النبات (سم) ، طول السنبله(سم) ، عدد ابراج السنبله ، عدد السنابل/م^٢ ، وزن ال ١٠٠٠ حبة (جم) ، محصول الحبوب ومحصول القش (كجم/فدان). علاوة علي ذلك كان صنف جميزة ٩ أفضل الأصناف في صفة التباين ل طرد السنابل مقارنة بالاصناف الأخرى.

- تأثر معنويًا كل من عدد الأيام ل طرد ٥٠% من السنابل ، مساحة ورقة العلم ، طول النبات ، طول السنبله ، عدد ابراج السنبله ، وعدد السنابل في المتر المربع ، محصول الحبوب والقش للفدان بالتفاعل بين معاملات الري و أصناف القمح.

- أدت معاملتي تحريم رية في طور اللبني و طور امتلاء الحبوب الي نقص في النسب المئوية لكل من الرطوبة، للدهون والكربوهيدرات في حبوب القمح. بينما ادت نفس المعاملتين الي زيادة في النسب المئوية لكل من البروتين ، الرماد والالياف الخام. أما بالنسبة للأصناف فقد سجل صنف جيزة ١٦٨ أقل قيم للنسب المئوية لكل من الرطوبة والكربوهيدرات وأيضا سجل نفس الصنف أعلى قيمة للنسبة المئوية للبروتين. بينما سجل صنف سدس ١٣ أعلى قيمة للنسبة المئوية للرماد ، سجل صنف سخا ٩٤ أعلى قيمة للنسبة المئوية للألياف وسجل الصنف مصر ١ أعلى قيمة للنسبة المئوية للدهون في حبوب القمح.

- أعطت معاملة التفاعل بين كل من تحريم رية في طور امتلاء الحبوب لصنف جيزة ١٦٨ أقل نسبة مئوية لكل من الرطوبة والكربوهيدرات وأيضاً أعطت نفس المعاملة أعلى نسبة المئوية للبروتين في الحبوب. بينما أعطت معاملة التفاعل بين كل من تحريم رية في الطور اللبني و طور امتلاء الحبوب لصنف سدس ١٣ أعلى قيم للنسبة المئوية لكل من للرماد والألياف علي الترتيب. وعند تحريم رية في الطور اللبني أو طور امتلاء الحبوب لصنف جيزة ١٦٨ أو صنف جيزة ٩ تم الحصول علي قيمة للنسبة المئوية للدهون في الحبوب.