

## Genetic evaluation and selection correlated response of growth traits in Japanese quail

M.M. El-Attrouny<sup>1#</sup>, E.A. Manaa<sup>2</sup> & S.I. Ramadan<sup>2</sup>

<sup>1</sup>Department of Animal Productions, Faculty of Agriculture at Moshtohor, Benha University, Egypt

<sup>2</sup>Animal and Poultry Production, Department of Animal Wealth Development, Faculty of Veterinary Medicine, Benha University, Moshtohor, Toukh 13736, Qalyubia, Egypt

(Received 6 December 2019; Accepted 13 April 2020; First published online 18 May 2020)

Copyright resides with the authors in terms of the Creative Commons Attribution 4.0 South African License.

See: <http://creativecommons.org/licenses/by/4.0/za>

Condition of use: The user may copy, distribute, transmit and adapt the work, but must recognize the authors and the South African Journal of Animal Science.

### Abstract

Objectives of the current study were to i) investigate effects of selection for bodyweight at four weeks old on bodyweight (BW) and bodyweight gain (BWG) across four generations; ii) estimate correlated response to selection for BW and BWG at different ages; and iii) document best linear unbiased prediction (BLUP) of genetic trends for BW and BWG across four generations of selection. A total of 3540 chicks from 444 sires and 885 dams were used to estimate heritabilities, and genetic and phenotypic correlations for growth traits, including BW at 0, 2, 4, and 6 weeks, and BWG between 0 and 2, 2 and 4, 4 and 6, and 0 and 6 weeks. The selection effects, correlated responses and genetic trend for BW and BWG across generations were quantified by applying the animal model. Estimates of heritability for BW and BWG ranged from 0.22 to 0.42 and from 0.18 to 0.23, respectively. Ranges of genetic and phenotypic correlations for BW varied from 0.31 to 0.92 and 0.05 to 0.65, respectively. Moreover, estimates of genetic and phenotypic correlations for BWG at different ages were from 0.12 to 0.72 and 0.17 to 0.60, respectively. Bodyweight and BWG estimates after four generations of selection were significantly higher than those of the base generation. Moreover, contrasts of generation means were significant across the four generations. The genetic trends across the generations clarified that BLUP estimates for BW and BWG gradually increased with the advance of generations until the fourth generation.

**Keywords:** best linear unbiased prediction, bodyweight, heritability, selection, genetic trend

<sup>#</sup>Corresponding author: Mahmoud.elatrouny@fagr.bu.edu.eg

### Introduction

Japanese quail constitute one of the most productive poultry species. They play an important role in the commercial poultry industry because of their high egg and meat production (Mizutani, 2003; Boni *et al.*, 2010). Moreover, they have many breeding advantages such as short generation interval, small floor space, high reproductive performance, early sexual maturity, rapid growth rate and high resistance to diseases (Baumgartner, 1994; Minvielle, 2004; Alkan *et al.*, 2010). All these characteristics made Japanese quail an excellent model for selection programmes. The poultry industry aims mainly to produce genetically superior animals with high productivity of egg and meat. Studies on selection programmes as the most important tool for achieving genetic gain in quail have been performed in two directions: the first was to improve egg production traits (Nestor *et al.*, 1983; Minvielle *et al.*, 2000; Durmus *et al.*, 2017); and the second to improve growth traits (Caron *et al.*, 1990; Marks, 1993; Balcioglu *et al.*, 2005; Ayatollahi Mehrgardi, 2013). In the second direction, weight gain was regarded as the most important selection criterion.

Selection for BW at four weeks in Japanese quail in long-term selection studies showed a positive correlation with BW in later weeks, genetic gain, age at sexual maturity and egg weight, and a negative correlation with reproductive traits and egg production rate (Marks, 1996; Khaldari *et al.*, 2010). Moreover, the phenotypic mass selection approach in long-term selection studies was followed by an increase in BW at four weeks old (Darden & Marks, 1988; Toelle *et al.*, 1991; Resende *et al.*, 2005; Varkoohi & Kaviani, 2014). This illustrates the quick response to selection for higher BW in Japanese quail.

Genetic evaluations of economic traits such as BW and BWG at different ages are of great interest in most poultry breeding programmes (Silva *et al.*, 2013). Therefore, most of the selection programmes have

been based on estimates of genetic parameters for growth traits (Nasiri Foomani *et al.*, 2014; Karami *et al.*, 2017). Several studies indicated that selection based on genetic parameters of growth traits resulted in good selection responses in BW of Japanese quail (Nestor *et al.*, 2000; Saatci *et al.*, 2006). In selection studies, several methodologies were used to estimate the selection response. One of these methods was regression analysis for the estimates of the breeding values on generations. This approach depends on the genetic parameters and the model. To the authors' knowledge, most selection studies overlooked the differences between genetic perspectives, such as selection-correlated responses, genetic trends and BLUP estimates.

The authors hypothesized that selection methods that depend on BLUP estimates and genetic trend will reveal promising improvement in growth performance of Japanese quail. Therefore, the main objectives of the current study were to investigate the selection effects, selection correlated responses, genetic trends and BLUP estimates for BW and BWG in Japanese quail through four selected generations.

## Materials and Methods

The current selection experiment was carried out at the poultry research farm at Faculty of Agriculture, Benha University, Egypt. Data of 3540 chicks that were obtained from 444 sires and 885 dams through five generations (base and four selected generations) were used in this study. Each breeding family consisted of one selected male and two selected females that were housed separately in a breeding cage (25 × 25 × 25 cm) with a sloping floor for collecting the pedigreed eggs. These eggs were incubated to get the first selected generation (G<sub>1</sub>), then the selection was practised further on G<sub>1</sub> to obtain the consecutive second (G<sub>2</sub>), third (G<sub>3</sub>) and fourth (G<sub>4</sub>) selected generations based on their BLUP estimates for BW at four weeks old. In each generation, the pedigree eggs in each breeding pen were collected daily for ten days and then incubated. The structures of the data collected during the experiment are presented in Table 1.

**Table 1** Numbers of sires, dams, females produced and chicks in a selection experiment with Japanese quail

Generation	Sires	Dams	Progeny (laying hens)	Chicks produced
0	125	249	260	688
1	94	188	270	781
2	82	164	264	684
3	72	144	272	687
4	71	140	266	700
Total	444	885	1332	3540

In each generation, newly hatched chicks were wing banded. Chicks were raised on floor cages until 12 days old. After that, chicks were transferred to battery brooders. At five weeks old, all females were transferred to individual laying cages for accurate pedigree identification. All birds were fed a standard diet containing 240 g/kg crude protein and 12.45 MJ/kg metabolizable energy during the rearing period according to NRC (1994). Feed and drinking water were offered to birds ad libitum. The lighting duration was 24 hours during the first week old, reduced to 23 hours from the second to four weeks old. In the laying house, the lighting regimen was set at 16 hours light and 8 hours dark.

Bodyweights of 3540 chicks were obtained in grams at 0, 2, 4 and 6 weeks old and presented as BW0, BW2, BW4, BW6, respectively. Bodyweight gain was calculated during the intervals from 0 to 2, 2 to 4, 4 to 6, and 0 to 6 weeks old and presented as BWG0-2, BWG2-4, BWG4-6, and BWG0-6, respectively. Variance-covariance components were estimated by REML using the computer package VCE6 (Groeneveld *et al.*, 2010). Multi-trait animal model (in matrix notation) was used to analyse growth traits using the following equation:

$$y = X\beta + Za + e$$

Where:  $y = nx1$  vector of observations with  $n =$  number of records,  
 $X =$  a  $n \times p$  incidence matrix which relates the data to a fixed effect of generation with five levels,

$\beta$  = a  $p \times 1$  vector of the fixed effects of generation,  
 $Z$  = an incidence matrix relating the data to additive genetic effects of the birds which are denoted by  $a$ , and  
 $e$  = vector of random residual effects.

Variance components for growth traits were computed using the package of VCE6. Heritability estimates ( $h_a^2$ ) were calculated from the results as:

$$h_a^2 = \frac{\sigma_a^2}{\sigma_a^2 + \sigma_e^2}$$

Where:  $\sigma_a^2$  and  $\sigma_e^2$  are estimates of the variances due to direct additive genetic effects and random error, respectively. The genetic ( $r_g$ ) and phenotypic ( $r_p$ ) correlations among growth traits were also estimated. Estimated covariances were obtained by solving the corresponding mixed model equations for bivariate models.

The best linear unbiased predictors of  $a$  for the birds with and without records were estimated using the PEST program (Groeneveld *et al.*, 2010). Genetic trend was measured by regression of the  $a$  values on generation number.

Estimates of correlated response in trait  $y$  to selection on trait  $x$  were calculated following Falconer and Mackay (1996):

$$CR_y = ih_x h_y r_g \sigma_{py}$$

Where:  $CR_y$  = the correlated response in trait  $y$ ,  
 $i$  = the selection intensity for trait  $x$ ,  
 $h_x$  and  $h_y$  = square roots of heritability for traits  $x$  and  $y$ , respectively,  
 $r_g$  = the genetic correlation of traits  $x$  and  $y$ , and  
 $\sigma_{py}$  = the phenotypic standard deviation of trait  $y$ .

### Results and Discussion

The actual means of BW were 9.5, 69.5, 152.2 and 241.4 g for chicks at 0, 2, 4 and 6 weeks old, respectively. These findings agree with the reported live BW of Japanese quail at various ages in previous studies (Sezer *et al.*, 2006; Varkoohi *et al.*, 2011; Silva *et al.*, 2013; Barbieri *et al.*, 2015). The means of BWG in Japanese quail were 4.3, 5.9, 6.4, and 5.5 g during the intervals from 0 to 2, 2 to 4, 4 to 6, and 0 to 6 weeks old, respectively. Coefficients of variation (CV%) ranged from 9.0% to 11.0% for BW and from 7% to 14% for BWG. Actual means, their standard deviations (SD) and percentages of variation (CV%) for BW0, BW2, BW4 and BW6 and BWG0-2, BWG2-4, and BWG4-6 and BWG0-6 weeks in Japanese quail are reported in Table 2.

**Table 2** Descriptive statistics of bodyweight and bodyweight gain for a selected population of Japanese quail

Traits	Number of records	Mean	Standard deviation	CV,%
Weight at hatching	3539	9.45	1.08	11
Weight at 2 weeks	3488	69.51	7.95	11
Weight at 4 weeks	3457	152.20	15.78	10
Weight at 6 weeks	3440	241.40	16.84	9
Gain hatching to 2 weeks	3488	4.29	0.54	13
Gain from 2 to 4 weeks	3457	5.91	0.83	14
Gain from 4 to 6 weeks	3440	6.37	0.77	12
Gain hatching to 6 weeks	3440	5.52	0.39	7

These current findings agree with those previous studies, which showed that CV% for BW ranged from 10.0 to 24.0 at various ages of Japanese quail (Narinc *et al.*, 2010; Varkoohi *et al.*, 2011; Barbieri *et al.*,

2015). Moreover, Özsoy and Aktan (2011) reported higher estimates of CV% for BW2 (20.7%) and BW4 (16.5%) in Japanese quail.

In this study, heritability estimates of BW were decreasing with the advancement of age of Japanese quail. Estimated  $h^2$  for BW0 (0.42) was approximately twofold the  $h^2$  of BW6 weeks (0.22). The current results agreed with those of previous studies (Shokoohmand *et al.*, 2007; Silva *et al.*, 2013; Ebrahimi *et al.*, 2018; Mohammadi-Tighsiah *et al.*, 2018). These findings could encourage the breeders to practise genetic selection at an early age in Japanese quail. In contrast, Saatci *et al.* (2003), Nasiri Foomani *et al.* (2014), and Karami *et al.* (2017) found that heritability values for BW of Japanese quail tended to increase with age. The estimated heritabilities ( $h^2$ ) and genetic and phenotypic correlations of BW and BWG of Japanese quail are shown in Table 3.

**Table 3** Estimates of heritability (on the diagonal), genetic correlation (above the diagonal) and phenotypic correlation (below the diagonal) and the associated standard errors for bodyweight and bodyweight gain of Japanese quail

	Bodyweight			
	Hatching	2 weeks	4 weeks	6 weeks
Hatching	0.42 ± 0.04	0.31 ± 0.07	0.44 ± 0.09	0.36 ± 0.12
2 weeks	0.15 ± 0.02	0.24 ± 0.04	0.59 ± 0.18	0.45 ± 0.16
4 weeks	0.08 ± 0.01	0.46 ± 0.13	0.36 ± 0.04	0.92 ± 0.20
6 weeks	0.05 ± 0.01	0.28 ± 0.07	0.65 ± 0.19	0.22 ± 0.04
	Bodyweight gain			
	Hatching to 2 weeks	2 to 4 weeks	4 to 6 weeks	Hatching to 6 weeks
Hatching to 2 weeks	0.19 ± 0.02	0.12 ± 0.03	0.33 ± 0.07	0.53 ± 0.05
2 to 4 weeks	0.18 ± 0.04	0.22 ± 0.02	0.28 ± 0.04	0.72 ± 0.15
4 to 6 weeks	0.17 ± 0.04	0.31 ± 0.07	0.18 ± 0.02	0.42 ± 0.09
Hatching to 6 weeks	0.27 ± 0.03	0.60 ± 0.11	0.48 ± 0.05	0.23 ± 0.02

Genetic and phenotypic correlations for BW at various ages (0, 2, 4, and 6 weeks) were positive and in the ranges of 0.31 to 0.92 and 0.05 to 0.65, respectively. For BW, estimates of genetic correlations between ages were higher than the corresponding phenotypic correlations. The highest value for genetic correlation was estimated between BW4 and BW6 (0.92) and the lowest was between BW0 and BW2 (0.31). Except for BW0 correlations with BW in various weeks, genetic correlations tended to increase with age. In addition, the correlation between consecutive closet BWs was higher than earlier. Genetic and phenotypic correlations were positive among BW at various ages, indicating that BW can be further improved with providing good management and climate conditions.

In agreement with the current findings, Silva *et al.* (2013) reported that genetic and phenotypic correlations between weekly BW of Japanese quail were 0.35 to 0.96 and 0.17 to 0.80, respectively. In contrast, Karami *et al.* (2017) revealed a higher estimated genetic correlation of 0.85, 0.90 and 0.91 between BW0 with BW7, BW14 and BW21, respectively, in Japanese quail through a random regression model.

Estimated  $h^2$  for BWG at different ages tended to be low, varying from 0.18 (BWG 4-6) to 0.23 (BWG 0-6). Estimates of genetic and phenotypic correlations for BWG at different ages (0 - 2, 2 - 4, 4 - 6, 0 - 6) were in the ranges of 0.12 to 0.72 and 0.17 to 0.60, respectively. The highest genetic correlation was estimated between BWG2-4 and BWG0-6 (0.72) and the lowest was 0.12 for earlier ages (between BWG0-2 and BWG2-4). Caetano *et al.*, (2017) reported that the estimate of  $h^2$  for BWG0-3 weeks in a meat-type quail population was 0.15. Also, in the study of Mohammadi-Tighsiah *et al.* (2018) the estimates of  $h^2$  for BWG0-5 weeks and BWG15-20 weeks were 0.08 and 0.10, respectively, which were lower than the present estimates in different periods.

Estimates of the selection for the four generations that were obtained by generalized least squares using VCE6 were superior ( $P < 0.05$ ) for most BW and BWG to the base population (Tables 4 and 5). The contrasts among estimates of these generations were significant ( $P < 0.05$ ). The fourth generation showed superiority in BW and BWG compared with the average of the first, second and third generations.

**Table 4** Generalized least-square estimates for bodyweights (g) in different generations and contrasts of the generation means illustrating pattern of selection response in Japanese quail

Generation means	Bodyweight			
	Hatching	2 weeks	4 weeks	6 weeks
0	8.81 ± 0.03 <sup>d</sup>	58.22 ± 1.21 <sup>e</sup>	134.47 ± 2.74 <sup>e</sup>	223.35 ± 3.12 <sup>e</sup>
1	9.37 ± 0.03 <sup>c</sup>	67.24 ± 1.26 <sup>d</sup>	150.77 ± 2.51 <sup>d</sup>	237.08 ± 3.42 <sup>d</sup>
2	9.70 ± 0.03 <sup>ab</sup>	72.58 ± 1.21 <sup>c</sup>	153.17 ± 2.74 <sup>c</sup>	243.97 ± 3.12 <sup>c</sup>
3	9.63 ± 0.03 <sup>b</sup>	74.57 ± 1.21 <sup>b</sup>	159.42 ± 2.74 <sup>b</sup>	250.13 ± 3.12 <sup>b</sup>
4	9.75 ± 0.03 <sup>a</sup>	75.16 ± 1.21 <sup>a</sup>	163.46 ± 2.74 <sup>a</sup>	253.13 ± 3.12 <sup>a</sup>
<i>Generation contrasts</i>				
0 vs 1	-0.56 ± 0.01 <sup>*</sup>	-9.02 ± 0.97 <sup>*</sup>	-16.30 ± 1.52 <sup>*</sup>	-13.73 ± 1.11 <sup>*</sup>
0 vs 2	-0.89 ± 0.01 <sup>*</sup>	-14.36 ± 1.03 <sup>*</sup>	-18.70 ± 1.58 <sup>*</sup>	-20.62 ± 1.23 <sup>*</sup>
0 vs 3	-0.82 ± 0.02 <sup>*</sup>	-16.35 ± 0.99 <sup>*</sup>	-24.95 ± 1.21 <sup>*</sup>	-26.78 ± 2.85 <sup>*</sup>
0 vs 4	-0.94 ± 0.04 <sup>*</sup>	-16.94 ± 1.01 <sup>*</sup>	-28.99 ± 2.52 <sup>*</sup>	-29.78 ± 2.63 <sup>*</sup>
1 vs 2	-0.33 ± 0.02 <sup>*</sup>	-5.34 ± 0.88 <sup>*</sup>	-2.40 ± 0.09 <sup>*</sup>	-6.89 ± 1.42 <sup>*</sup>
1 vs 3	-0.26 ± 0.02 <sup>*</sup>	-7.33 ± 0.91 <sup>*</sup>	-8.65 ± 0.87 <sup>*</sup>	-13.05 ± 2.14 <sup>*</sup>
1 vs 4	-0.38 ± 0.01 <sup>*</sup>	-7.92 ± 0.91 <sup>*</sup>	-12.69 ± 1.11 <sup>*</sup>	-16.05 ± 1.21 <sup>*</sup>
2 vs 3	0.07 ± 0.01	-1.99 ± 0.98 <sup>*</sup>	-6.25 ± 0.58 <sup>*</sup>	-6.16 ± 1.52 <sup>*</sup>
2 vs 4	0.07 ± 0.01	-2.75 ± 0.89 <sup>*</sup>	-10.29 ± 1.12 <sup>*</sup>	-9.16 ± 1.91 <sup>*</sup>
3 vs 4	-0.12 ± 0.01 <sup>*</sup>	-0.59 ± 0.80 <sup>*</sup>	-4.04 ± 0.88 <sup>*</sup>	-3.0 ± 1.41 <sup>*</sup>

<sup>a,b,c,d</sup> Within a column, estimates with a common superscript do not differ at  $P=0.05$

\* Contrast is significantly different from 0 at  $\alpha=0.05$

**Table 5** Generalized least-square estimates for bodyweight gain (g/d) in different generations and contrasts of the generation means illustrating the pattern of selection response in Japanese quail


Generation means	Hatching to 2 weeks	Weeks 2 to 4	Weeks 4 to 6	Hatching to 6 weeks
0	3.52 ± 0.10 <sup>d</sup>	5.44 ± 0.19 <sup>e</sup>	6.34 ± 0.24 <sup>b</sup>	5.10 ± 0.17 <sup>e</sup>
1	4.13 ± 0.13 <sup>c</sup>	5.96 ± 0.22 <sup>c</sup>	6.16 ± 0.25 <sup>c</sup>	5.42 ± 0.19 <sup>d</sup>
2	4.49 ± 0.11 <sup>b</sup>	5.75 ± 0.21 <sup>d</sup>	6.48 ± 0.24 <sup>a</sup>	5.57 ± 0.17 <sup>c</sup>
3	4.63 ± 0.11 <sup>a</sup>	6.06 ± 0.21 <sup>b</sup>	6.47 ± 0.24 <sup>a</sup>	5.72 ± 0.17 <sup>b</sup>
4	4.67 ± 0.11 <sup>a</sup>	6.36 ± 0.21 <sup>a</sup>	6.40 ± 0.24 <sup>ab</sup>	5.79 ± 0.17 <sup>a</sup>
<i>Generation contrasts</i>				
0 vs 1	-0.61 ± 0.08 <sup>*</sup>	-0.52 ± 0.08 <sup>*</sup>	0.18 ± 0.07 <sup>*</sup>	-0.32 ± 0.07 <sup>*</sup>
0 vs 2	-0.97 ± 0.08 <sup>*</sup>	-0.31 ± 0.09 <sup>*</sup>	-0.14 ± 0.08 <sup>*</sup>	-0.47 ± 0.09 <sup>*</sup>
0 vs 3	-1.11 ± 0.08 <sup>*</sup>	-0.62 ± 0.08 <sup>*</sup>	-0.13 ± 0.07 <sup>*</sup>	-0.62 ± 1.01 <sup>*</sup>
0 vs 4	-1.15 ± 0.10 <sup>*</sup>	-0.92 ± 0.09 <sup>*</sup>	-0.06 ± 0.09	-0.69 ± 1.01 <sup>*</sup>
1 vs 2	-0.36 ± 0.07 <sup>*</sup>	0.21 ± 0.08 <sup>*</sup>	-0.32 ± 0.08 <sup>*</sup>	-0.15 ± 0.09 <sup>*</sup>
1 vs 3	-0.50 ± 0.08 <sup>*</sup>	-0.10 ± 0.07 <sup>*</sup>	-0.31 ± 0.06 <sup>*</sup>	-0.30 ± 0.08 <sup>*</sup>
1 vs 4	-0.54 ± 0.05 <sup>*</sup>	-0.40 ± 0.09 <sup>*</sup>	-0.24 ± 0.07 <sup>*</sup>	-0.37 ± 1.11 <sup>*</sup>
2 vs 3	-0.14 ± 0.07 <sup>*</sup>	-0.31 ± 0.09 <sup>*</sup>	0.10 ± 0.07	-0.15 ± 0.09 <sup>*</sup>
2 vs 4	-0.18 ± 0.08 <sup>*</sup>	-0.61 ± 0.07 <sup>*</sup>	0.08 ± 0.07	-0.22 ± 0.09 <sup>*</sup>
3 vs 4	-0.04 ± 0.09	-0.30 ± 0.08 <sup>*</sup>	0.07 ± 0.08	-0.07 ± 0.09

<sup>a,b,c,d</sup> Within a column, estimates with a common superscript do not differ at  $P=0.05$

\* Contrast is significantly different from 0 at  $\alpha=0.05$

Khaldari *et al.* (2010) reported that the mean of BW at four weeks old in the S line and C line in the last generation (fourth generation) was 200.5 and 170.9 g, respectively, a 17.3% cumulative genetic improvement, or 5.8% improvement per generation. Magda *et al.* (2010) showed that the means of six weeks' BW was 172.2, 186.3, 202.9, and 214.7 g for the base, first, second and third generations.

The BLUP estimates for BW0, BW2, BW4, and BW6 increased gradually as the generation advanced until they reached 0.07, 1.17, 3.90, and 6.60 g in G<sub>4</sub>, respectively (Figure 1a, b, c, and d). A similar genetic trend for BWG was recorded where the BLUP estimates for BWG0-2, BWG2-4, BWG4-6, and BWG0-6 increased gradually as the generation advanced until they reached 0.05, 0.22, 0.25, and 0.18 g in G<sub>4</sub>, respectively (Figure 1e, f, g, and h).



**Figure 1** Genetic trend for growth traits of bodyweight and bodyweight gain across the base and selected generations

This increased genetic trend in the selected generations illustrated the efficiency of selection for BW at four weeks old. The estimation of genetic trend was necessary for the assessment of the breeding programmes. The current results agreed with those of Hussain *et al.* (2014), who observed a linear increasing trend of BW at three weeks old in selected populations as the generations progressed. On the other hand, Narinc *et al.* (2016) found that the mean of estimated BLUP values for BW in base population, first and second selected generations flocks were found to be 0.13, 0.20, and 0.34, respectively.

The correlated response for unselected traits showed a positive correlation for increasing all BW at ages (Table 6). In all BW stages, the improve in BW decreased with the advance of generations. However, the cumulative correlated response of selection for BW0, BW2 and BW6 was increased dramatically by 0.83, 10.88 and 25.45 g, respectively. A similar trend was observed for BWG at different age intervals. These findings agree with the results of previous studies (Anthony *et al.*, 1996; Ayatollahi Mehrgardi, 2013; Varkoohi, 2014; Hussien *et al.*, 2016). These responses indicate that selection for BW at four weeks was associated with an increase in BW at different ages. The authors found that the correlated responses were due to the high genetic correlation between BW at four weeks old and other ages. This revealed that selection for higher BW at an early age would bring a concomitant improvement in BW at later ages as a correlated response to the selection.

**Table 6** Correlated response to selection for four-week bodyweight in Japanese quail over four generations

	Generation number				Cumulative
	1	2	3	4	
<i>Bodyweight</i>					
Hatching	0.44	0.27	0.04	0.08	0.83
2 weeks	5.42	3.54	1.41	0.51	10.88
6 weeks	12.32	4.93	5.35	2.85	25.45
<i>Bodyweight gain</i>					
Hatching to 2 weeks	0.54	0.31	0.11	0.06	1.06
Weeks 2 to 4	0.42	0.21	0.22	0.27	1.12
Weeks 4 to 6	0.14	0.29	0.08	0.05	0.56
Weeks 0 to 6	0.29	0.17	0.13	0.04	0.63

## Conclusion

Japanese quail respond quickly to selection for BW at four weeks old. The selected generations indicate that selection for BW at four weeks old was associated with an improvement in BW and BWG at other ages. Additionally, cumulative correlated responses of selection, genetic trends and BLUP estimates for BW and BWG increased gradually until the fourth generation. Hence, genetic improvement in growth performance could be achieved through practising BW selection at four weeks old and more improvement could be achieved through selection for more generations based on BLUP estimates.

## Acknowledgments

The authors are grateful to Department of Animal Production, Faculty of Agriculture in Benha University, Egypt, for the facilities during the breeding experiment of the quail. This work was supported by the research project entitled Genetic Improvement for Japanese Quail Using Molecular Marker Tools from the Scientific Research Fund (SRF), Benha University, Egypt.

## Authors' Contributions

MME and EAM designed the experiment and collected data. MME and SIR analysed the data and interpreted the results. MME wrote the draft manuscript and all authors cooperated in revising and finalizing the manuscript.

## Conflict of Interest Declaration

The authors have declared that there are no competing interests.

## References

- Alkan, S., Karabağ, K., Galiç, A., Karslı, T. & Balçioğlu, M.S., 2010. Determination of bodyweight and some carcass traits in Japanese quails (*Coturnix coturnix japonica*) of different lines. *Kafkas Univ. Vet. Fak. Derg.* 16:277-280. DOI: 10.9775/kvfd.2009.687
- Anthony, N.B., Nestor, K. E. & Marks, H.L., 1996. Short-term selection for four-week bodyweight in Japanese quail. *Poult. Sci.* 75, 1192-1197.
- Ayatollahi Mehrgardi, A., 2013. Divergent selection for four-week bodyweight in Japanese quail (*Coturnix coturnix japonica*): Response to selection and realized heritability. *J. Livest. Sci. Technol.* 1, 57-59.
- Balçioğlu, M.S., Kızılkaya, K., Yolcu, H.I. & Karabağ, K., 2005. Analysis of growth characteristics in short-term divergently selected Japanese quail. *S. Afr. J. Anim. Sci.* 35, 83-89.
- Barbieri, A., Ono, R.K., Cursino, L.L., Farah, M.M., Pires, M.P. Bertipaglia, T.S., Pires, A.V., Cavani, L., Carreño, L.O.D. & Fonseca, R., 2015. Genetic parameters for bodyweight in meat quail. *Poult. Sci.* 94, 169-171. <https://doi.org/10.3382/ps/peu062>
- Baumgartner, J., 1994. Japanese quail production, breeding and genetics. *Worlds Poult. Sci. J.* 50227-235.
- Boni, I., Nurul, H. & Noryati, I., 2010. Comparison of meat quality characteristics between young and spent quails. *Int. Food Res. J.* 17, 661-667.
- Caetano, G.C., Mota, R.R., da Silva, D.A., de Oliveira, H.R., Viana, J.M.S., de Siqueira, O.H.G., Freitas, P.H.F. & de Silva, F.F., 2017. Bayesian estimation of genetic parameters for individual feed conversion and bodyweight gain in meat quail. *Livest. Sci.* 200, 76-79. <https://doi.org/10.1016/j.livsci.2017.04.011>
- Caron, N., Minvielle, F., Desmarais, M. & Poste, L.M., 1990. Mass selection for 45-day bodyweight in Japanese quail: selection response, carcass composition, cooking properties, and sensory characteristics. *Poult. Sci.* 69:1037-1045. <http://dx.doi.org/10.3382/ps.0691037>
- Darden, J.R. & Marks, H.L. 1988. Divergent selection for growth in Japanese quail under split and complete nutritional environments. 1. Genetic and correlated responses to selection. *Poult. Sci.* 67, 519-529.
- Durmus, I., Alkan, S., Narinc, D., Karabag, K. & Karsli, T., 2017. Effects of mass selection on egg production on some reproductive traits in Japanese quail. *Europ. Poult. Sci.* 1, 81.
- Ebrahimi, K., Dashab, G.R., Faraji-Arough, H. & Rokouei, M., 2019. Estimation of additive and non-additive genetic variances of bodyweight in crossbreed populations of the Japanese quail. *Poult. Sci.* 98, 46-55. <https://doi.org/10.3382/ps/pey357>
- Falconer, D.S. & Mackay, F.C., 1996. Introduction to quantitative genetics. Fourth edition. Longman Group, Harlow, Essex, England. Pp.108-183.
- Groeneveld, E., Kovac, M. & Mielenz, N., 2010. VCE – user's guide and reference manual, version 6.0. Institute of Farm Animal Genetics, Neustadt, Germany.
- Hussain, J., Akram, M., Sahota, A.W., Javed, K., Ahmad, H.A., Mehmood, S., Jatoi, A.S. & Ahmad, S., 2014. Selection for higher three-week bodyweight in Japanese quail: 2. Estimation of genetic parameters. *J. Anim. Plant. Sci.* 24, 869-873.
- Hussen, S.H., Abdulrahman Al-Khdri, A.M. & Hassan, A.M., 2016. Response to selection for bodyweight in Japanese quail (*Coturnix coturnix japonica*). *Iran. J. Appl. Anim. Sci.* 453-459.
- Karami, K., Zerehdaran, S., Tahmoorespur, M., Barzanoi, B. & Lotfi, E., 2017. Genetic evaluation of weekly bodyweight in Japanese quail using random regression models. *Br. Poult. Sci.* 58,13-18. <https://doi.org/10.1080/00071668.2016.1236362>
- Khaldari, M., Pakdel, A., Mehrabani Yegane, H., Nejati Javaremi, A. & Berg, P., 2010. Response to selection and genetic parameters of body and carcass weights in Japanese quail selected for 4-week bodyweight. *Poult. Sci.* 89, 1834-1841. <https://doi.org/10.3382/ps.2010-00725>
- Magda, I., Abo Samaha, M., Sharaf, M. & Hemeda, S.A., 2010. Phenotypic and genetic estimates of some productive and reproductive traits in Japanese quails. *Egypt. Poult. Sci.* 30, 875-892.
- Marks, H.L., 1993. Carcass composition, feed intake, and feed efficiency following long-term selection for four-week bodyweight in Japanese quail. *Poult. Sci.* 72:1005-1011. <https://doi.org/10.3382/ps.0721005>
- Marks, H.L., 1996. Long-term selection for bodyweight in Japanese quail under different environments. *Poult. Sci.* 75, 1198-1203. <https://doi.org/10.3382/ps.0751198>
- Minvielle, F., Monvoisin, J.L., Costa, J. & Maeda, Y., 2000. Long-term egg production and heterosis in quail lines after within-line or reciprocal recurrent selection for high early egg production. *Br. Poult. Sci.* 41, 150-157. <https://doi.org/10.1080/713654914>
- Minvielle, F., 2004. The future of Japanese quail for research and production. *Worlds. Poult. Sci. J.* 60, 500-507. <https://doi.org/10.1079/WPS200433>
- Mizutani, M., 2003. The Japanese quail. Laboratory Animal Research Station, Nippon Institute for Biological Science, Kobuchizawa, Yamanashi, Japan, 408-0041.
- Mohammadi-Tighsiah, A., Maghsoudi, A., Bagherzadeh-Kasmani, F., Rokouei, M. & Faraji-Arough, H., 2018. Bayesian analysis of genetic parameters for early growth traits and humoral immune responses in Japanese quail. *Livest. Sci.* 216, 197-202. <https://doi.org/10.1016/j.livsci.2018.07.012>
- Narinc, D., Aksoy, T. & Karaman, E., 2010. Genetic parameters of growth curve parameters and weekly bodyweights in Japanese quails (*Coturnix coturnix japonica*). *J. Anim. Vet. Adv.* 9, 501-507.
- Narinc, D., Aksoy, T. & Kaplan, S., 2016. Effects of multi-trait selection on phenotypic and genetic changes in Japanese quail (*Coturnix coturnix japonica*). *J. Poult. Sci.* 53, 103-110, DOI: 10.2141/jpsa.0150068
- Nasiri Foomani, N., Zerehdaran, S., Ahani Azari, M. & Lotfi, E., 2014. Genetic parameters for feed efficiency and bodyweight traits in Japanese quail. *Br. Poult. Sci.* 55, 298-304. <https://doi.org/10.1080/00071668.2014.925088>



- Nestor, K.E., Anderson, J.W. & Patterson, R.A., 2000. Effects of selection for increased bodyweight, egg production, and shank width on developmental stability in turkeys. *Poult. Sci.* 79, 937-945. <https://doi.org/10.1093/ps/79.7.937>
- Nestor, K.E., Bacon, W.L. & Lambio, A.L., 1983. Divergent selection for egg production in *Coturnix coturnix japonica*. *Poult. Sci.* 62, 1548-1552. <https://doi.org/10.3382/ps.0621548>
- NRC, 1994. Nutrient requirements of poultry. 9th revised edition. National Academic Press, Washington, DC, USA.
- Özsoy, A.N. & Aktan, S., 2011. Estimation of genetic parameters for body weight and egg weight traits in Japanese quails. *Trends Anim. Vet. Sci.* 2(1), 17-20.
- Resende, R.O., Martins, E.N., Georg, P.C., Paiva, E., Conti, A.C.M., Santos, A.I., Sakaguti, E.S. & Murakami, A.E., 2005. Variance components for bodyweight in Japanese quails (*Coturnix japonica*). *Braz. J. Poult. Sci.* 7, 23-25. <https://doi.org/10.1590/S1516-635X2005000100004>
- Saatci, M., Ap Dewi, I. & Aksoy, A.R., 2003. Application of REML procedure to estimate the genetic parameters of weekly live weights in one to one sire and dam pedigree recorded Japanese quail. *J. Anim. Breed. Genet.* 120, 23-28. <https://doi.org/10.1046/j.1439-0388.2003.00370.x>
- Saatci, M., Omed, H. & Ap Dewi, I., 2006. Genetic parameters from univariate and bivariate analyses of egg and weight traits in Japanese quail. *Poult. Sci.* 85, 185-190. <https://doi.org/10.1093/ps/85.2.185>
- Sezer, M., Berberoglu, E. & Ulutas, Z., 2006. Genetic association between sexual maturity and weekly live-weights in laying-type Japanese quail. *S. Afr. J. Anim. Sci.* 36, 142-148. <https://doi.org/10.4314/sajas.v36i2.3997>
- Shokoohmand, M., Kashan, N.E.J. & Maybody, M. E., 2007. Estimation of heritability and genetic correlations of bodyweight in different age for three strains of Japanese quail. *Int. J. Agric. Biol.* 9, 945-947.
- Silva, L.P., Ribeiro, J.C., Crispim, A.C., Silva, F.G., Bonafé, C.M., Silva, F.F. & Torres, R.A., 2013. Genetic parameters of bodyweight and egg traits in meat-type quail. *Livest. Sci.* 153, 27-32. <https://doi.org/10.1016/j.livsci.2013.01.014>
- Toelle, V.D., Havenstein, G.B., Nestor, K.E. & Harvey, W.R., 1991. Genetic and phenotypic relationships in Japanese quail. 1. Bodyweight, carcass, and organ measurements. *Poult. Sci.* 70, 1679-1688. <https://doi.org/10.3382/ps.0701679>
- Varkoohi, S. & Kaviani, K., 2014. Genetic improvement for bodyweight of Japanese quail. *Annu. Res. Rev. Biol.* 347-353. <https://doi.org/10.9734/ARRB/2014/5281>
- Varkoohi, S., Pakdel, A., Moradi Shahr Babak, M., Nejati Javaremi, A., Kause, A. & Zaghari, M., 2011. Genetic parameters for feed utilization traits in Japanese quail. *Poult. Sci.* 90, 42-47. DOI: 10.3382/ps.2010-01072