



---

## CORRELATED RESPONSE FOR EGG QUALITY TRAITS AFTER THREE GENERATIONS OF SELECTION FOR EGG NUMBER IN BENHA CHICKENS

El-Attrouny, M.M.<sup>1</sup>, Iraqi, M.M.<sup>1</sup>, Khalil, M.H.<sup>1</sup> and El-Moghazy, Gihan M.<sup>2</sup>

<sup>1</sup>Dep. of Anim. Prod., Fac. of Agric. at Moshtohor, Benha Uni., Egypt.

<sup>2</sup>Regional Centre for Food and Feed, Agri. Res. Center, Egypt.

**Corresponding author:** Iraqi, M.M Email: [mahmoud.iraqi@fagr.bu.edu.eg](mailto:mahmoud.iraqi@fagr.bu.edu.eg)

Received: 02/09/2021

Accepted: 28 /09/2021

---

**ABSTRACT:** The current study investigated the effect of selection for increased egg number during the 1<sup>st</sup> 90 days of laying on egg quality traits over four generations in Benha chickens. A total of 18 cockerels and 180 pullets were selected according to their BLUP values for egg number at the first 90 days of laying from Benha base population (control). Data of 4242 eggs produced by 756 pedigreed hens fathered by 69 sires and mothered 484 dams from four generations (base and three selected generations) were used to estimate heritability, genetic and phenotypic correlations for egg quality traits (Egg weight (EW), Albumen weight (AW),Yolk weight (YW), Shell weight (SW), Haugh unit (HU), Egg shape index (ESI), Albumen index (AI), Yolk index (YI) and Egg shell thickness (EST)). The selection effects, correlated responses and the genetic and phenotypic trends for egg quality traits across generations were quantified or clarified applying the updated approach of the animal model program of BLUPF90. Heritability estimates were moderate or low for all traits ( $h^2 = 0.05$  to  $0.47$ ). Also, genetic correlations for EW was closely positively genetically correlated with AW, YW, SW, with magnitudes of 0.99, 0.57, 0.69, respectively. The three selected generations were superior ( $P < 0.05$ ) in most egg quality traits than the base generation and the contrasts among estimates of these generations were significant ( $P < 0.05$ ). The phenotypic and genetic trend increased for all egg quality traits. While genetic trend decreased for AW. The accumulative correlated selection responses were 1.18 g for EW, 0.31g for SW, 5.27 for HU, and 0.06 mm for EST. Conclusion, these findings stated superiorities of selected generations indicating selection for egg number during the first 90 days of laying in Benha chickens was associated with an improvement in egg quality traits.

---

**Keywords:** selection, egg quality traits, genetic and phenotypic trends, correlated responses.

---

## INTRODUCTION

Egg quality is one of the most important requirements of today's market to guarantee the integrity of the egg and to reduce the numbers of eggs lost on the way to the consumer. According to Harms *et al.* (1996), approximately 6 to 8% of the total egg production is not usable or marketable due to the poor quality of shells. Hunton (1995) observed major financial losses during routine handling and transport from producer to retail outlets. Therefore, eggshell stability traits play a major role because only eggs with an intact shell are considered salable. Thus, if egg quality, and specifically, eggshell stability, is guaranteed, the layer industry could increase the number of salable eggs produced by each hen housed.

Genetic selection helps to continuously improve the laying performance, efficiency, livability, and adaptability of the birds to different environments as well as egg quality (Preisinger and Flock, 2000). For this purpose, more than 30 different traits are included in the selection index of the laying hen breeding programs. All traits are closely monitored and their relations to each other are considered, making it possible for a balanced genetic progress and improvements in different traits even when they are negatively correlated with each other. Estimations of genetic parameters for different egg quality traits have been reported in literature (Dunn *et al.*, 2005; Icken *et al.*, 2006; Flock *et al.*, 2007; El-Attrouny and Iraqi, 2021). According to Dunn *et al.* (2005) and Flock *et al.* (2007), heritabilities ( $h^2$ ) are ranged from 0.52 to 0.64 for EW and vary from  $h^2 = 0.33$  to 0.53 for Kdyn in brown and white egg layers. Egg weight (EW) is a major

criterion in determining the market price system, and according to Flock *et al.* (2007), EW is easily adaptable to the requirements of the market thanks to its high heritability. The main objectives of the present study were: 1) Estimate the genetic parameters and BLUP estimates across three generations of selection for egg quality traits, 2) Effect of selection quantify and comparing the generation contrasts, 3) clarify the phenotypic and genetic trends across generations, and 4) Estimate the correlated responses in egg quality over the three generations of selection in Benha chickens.

## MATERIALS AND METHODS

### Breeding plane and management:

A selection experiment was carried out at the Poultry Research Farm, Department of Animal Production, Faculty of Agriculture, Benha University, Egypt during the period from 2011 and terminated in 2015. A synthesized line of chickens, Benha line (Bl) was used in this study. This line has been developed from crossing Golden Montazah with White Leghorns chickens and using the BLUP estimates to select the birds for three generations (Iraqi *et al.*, 2013).

A total of 18 cockerels and 180 pullets were selected according to their BLUP estimates for egg number during the first 90 days of egg production from Benha line base population. Data of 4242 egg collected on 756 hens produced by 69 sires and 484 dams from four generations (base and three selected generations). Egg number (EN) was recorded for each hen by counting the number of eggs laid during the first 90 days after sexual maturity. Each selected cock was mated with 10 selected hens and housed separately in breeding pen to produce the first generation of selection ( $G_1$ ), consequently selection was practiced

## **selection, egg quality traits, genetic and phenotypic trends, correlated responses.**

further for two generations to produce the 2<sup>nd</sup> (G<sub>2</sub>) and 3<sup>rd</sup> (G<sub>3</sub>) generations of selection. The pedigreed eggs from each individual breeding pen for the three selection generations (G<sub>1</sub>, G<sub>2</sub> and G<sub>3</sub>) were collected daily for 15 days and then incubated (El-Attrouny *et al.*, 2019). The structures of data collected from all generations are presented in Table 1.

Upon hatch, chicks produced from all generations were wing - banded and reared in floor brooder, then transferred to the floor pens. Chicks produced from all generations were fed *ad libitum* during growing (from hatch up to 8 weeks of age), rearing (from 8-20 weeks of age) and laying (more than 20 weeks of age) periods on diets containing 21% protein and 2950 kcal/kg, 18% protein and 2850 kcal/kg, and 16% protein and 2700 kcal/kg, respectively. The feed requirements were supplied according to NRC (1994). All the birds were treated and medicated similarly throughout the experimental period.

### **Data and model of analysis**

Records on 4242 eggs from different generations were collected to study egg quality traits. Which were egg weight (EW), albumen weight (AW), yolk weight (YW), shell weight (SW), Haugh unit (HU), egg shape index (ESI), albumen index (AI), yolk index (YI) and shell thickness (EST).

Multi-trait animal model (in matrix notation) used to analyze egg quality traits (as characters of the hen) was as follows:

$$y = Xb + Z_a u_a + Z_p u_p + e$$

Where:  $y = n \times 1$  vector of observation of the hen,  $n =$  number of records;  $X =$  design matrix of order  $n \times p$ , which is related to the fixed effects of generation (four levels), month-year (12 levels);  $b = p \times 1$  vector of the fixed effects of

generation and month-year  $Z_a =$  the incidence matrix relating records to the additive genetic effect of the hen;  $u_a =$  the vector of random additive genetic of the hen;  $Z_p =$  the incidence matrix relating records to random permanent environmental effect of the hen;  $u_p =$  the vector of random permanent environmental effect of the hen; and  $e = n \times 1$  vector of random residual effects, NID (0,  $\sigma^2 e$ ).

Heritability estimates of egg quality traits

were computed as:

$$h^2_a = \frac{\sigma_a^2}{\sigma_a^2 + \sigma_p^2 + \sigma_e^2}$$

Where:  $s_a^2$ ,  $s_p^2$  and  $s_e^2$  are variances due to the effects of direct additive genetic, permanent environment and random error, respectively. The genetic ( $r_g$ ) and phenotypic ( $r_p$ ) correlations between egg quality traits were also estimated.

The BLUP were estimated using BLUPF90 Fortran program (Misztal *et al.*, 2014). Bird solutions are computed from the pedigree file, one bird at a time (for both birds with and without records, i.e. hens, sires and dams). These BLUP estimates were calculated based on the theory of Kennedy (1989).

The estimated (co) variances were used to estimate the fixed and the random effects by solving the corresponding mixed model equations using the BLUPF90 software (Misztal *et al.*, 2014). The estimates of the error (co) variance matrix were also obtained. Estimates of the differences between the four generations of selection were obtained by generalized least-squares analysis using the BLUPF90 software.

The phenotypic trend was measured as the regression of least squares estimates on generation number, while the genetic trend was measured by regressing the BLUP estimates on generation number. As stated before, the BLUP estimates of

the birds with and without records were estimated using the BLUPF90 program (Misztal *et al.*, 2014).

The correlated selection response ( $CR_Y$ ) in each trait (Y) was calculated using the following equation of Falconar and Meckay (1996):  $CR_Y = (i_X) (h_X) (h_Y) (r_{GXY}) (\sigma_{PY})$ ; where:  $i_X$  is the selection intensity assuming to be one for comparison only,  $h_X$  and  $h_Y$  are the square roots of heritability estimates of the X and Y traits, respectively,  $r_{GXY}$  is the genetic correlation between the two traits and  $\sigma_{PY}$  is the standard deviation of phenotypic value of trait Y.

## RESULTS AND DISCUSSION

### Actual means and variations:

The actual means for egg components, ESI and EST in Benha chickens were estimated and illustrated in Table 2. The means for EW, AW, YW and SW were 48.3, 27.6, 14.6 and 6.1 g, respectively. El-Garhy (2004) found that the means for EW, AW, YW, SW and EST were 43, 24, 13.4, 5.6 g and 0.31 mm in chickens. Khawja *et al.* (2013) found that the actual means of EW, AW and YW were 43.4, 22 and 14 g in Fayoumi chickens. Khalil *et al.* (2013) showed that the averages of EW, AW, YW and SW were 44.0, 24.2, 14.4 and 5.5 g, respectively in Golden Montazah chickens. Taha and Abd El-Ghany (2013) reported that means of EW in Mandarah and EL-Salam strains were 50.6 and 52.5 g, respectively. The actual means of HU, ESI, AI, YI and EST 90.6, 78.4, 8.0, 44.0 % and 0.33 mm, respectively (Table 2). The shell thickness means was 0.33 mm. Abd El-Latif (2001) reported that the means for HU was 69.2 and 68.4 in Dandarawi and Golden Montazah chicken, respectively. Iraqi (2002) reported that the means of ESI, AI and YI were 77.5, 11.3 and 47.83 %, respectively in Mandara chickens, while

they were 77.2, 47.5 and 86.6 % for the same traits in Matrouh chickens, respectively.

The percentages of variation ranged from 9.8 to 28.6 % for egg component traits and from 0.11 to 0.21 % for shape index traits (Table 2) and the variation for shell thickness was 0.12 %. Olawumi and Ogunlade (2008) found that the percentage of variation ranged from 7.8 to 9.8 % for egg components and from 3.7 to 9.0 % for shape index traits and 12.6 % for EST in Brown layer breeds. Olawumi and Jobin (2013) reported that the percentages of variation were 9.6, 22.1, 33.0, 11.4, 7.3, 7.8 and 5.4 % for EW, AW, YW, SW, EST, HU, ESI, respectively in Rhode Island White chickens.

### Variance components and heritability:

Estimates of additive ( $\sigma_a^2$ ), permanent ( $\sigma_{PE}^2$ ) and residual ( $\sigma_e^2$ ) variances for egg quality traits are given in Table (3). The estimates of  $\sigma_a^2$  were low and intermediate for egg components traits (ranged from 0.75 to 16.1g) compared to shape index trait (ranged from 0.07 to 1.7 %) and EST (2.1 mm). These results indicate that egg components traits had high additive genetic variance comparable to shape indexes and shell thickness traits. Therefore, improving egg components traits by selection could be possible. These estimates of additive genetic variance are agreed with Iraqi (2002) using an animal model, while they are disagreed with those reported by Abou El-Ghar *et al.* (2011) and El-Attrouny (2011).

Egg components traits had intermediate estimate of  $h^2$ , ranging from 0.17 to 0.47 for egg components, 0.06 and 0.25 for shape indexes and 0.23 for shell thickness (Table 3). It seems that egg components are largely influenced by genes of

## **selection, egg quality traits, genetic and phenotypic trends, correlated responses.**

additive effect and thus these traits could be improved by selection. Based on multi-trait animal model, Iraqi (2002) found that heritability estimates were 0.67, 0.50, 0.30 and 0.25 for EW, AW, YW and SW, respectively in Mandarah and Matrouh chickens. Zhang *et al.* (2005) found that estimates of  $h^2$  for EW, AW, YW, SW, HU and ESI in brown-egg dwarf layers were 0.63, 0.59, 0.45, 0.64, 0.41 and 0.34, respectively. Dasari *et al.* (2013) reported that the estimates of  $h^2$  ranged from 0.12 to 0.32 for egg components, from 0.10 to 0.23 for shape indexes, from 0.27 to 0.41 for shell thickness in different strains of White Leghorns. On the other hand, Younis *et al.* (2014) found that estimates of  $h^2$  for ESI, AI, AI, HU and EST were 0.39, 0.29, 0.46, 0.18 and 0.21, respectively in Dokki\_4 chickens. Ashour *et al.* (2015) found that estimate of  $h^2$  for ESI, AI, YI, EST and HU were 0.45, 0.36, 0.18, 0.07 and 0.15, respectively in El-Salam chickens. Also, Rath *et al.* (2015) reported that the estimates of  $h^2$  for egg components ranged from 0.14 to 0.57, from 0.40 to 0.59 for shape indexes and 0.44 for shell thickness in White Leghorn.

### **Genetic and phenotypic correlation**

Estimated of  $r_G$  were positive and closely correlated between EW & AW (0.99), EW & YW (0.57) and EW & SW (0.69) are presented in Table (4). It seems that pullets giving high egg weight could have higher albumin, yolk and shell weights. Also, positive  $r_G$  between AW&AI (0.59) and YW&YI (0.61) were recorded. Hanafi and El-Labban (1990) found that highly positive genetic correlation between EW & AW (0.69) and EW&SW (0.52) in Dokki-4 chickens. Zhang *et al.* (2005) found that  $r_G$  recorded among EW and each of AW, YW, SW ranged from 0.67 to 0.97 in brown-egg layer. El-

Attrouny (2011) found that highly positive  $r_G$  among EW and each of AW (0.80), YW (0.70) and SW (0.53) in Golden Montazah and White Leghorn and their crosses.

Estimates of  $r_p$  presented in Table ( 4 ), the  $r_G$  were positive and highly correlated between EW & AW, EW & YW and EW&SW; being 0.85, 0.63 and 0.54, respectively, while the correlations between AW&AI (0.02) and YW & YI (-0.26) were low and negative in Benha chickens. Udoh *et al.* (2012) indicted that correlations between EW & AW were positive and ranged from 0.68 to 0.79 and from 0.56 to 0.77 between EW & YW in three Nigerian chickens. Dasari *et al.* (2013) found that the estimates of  $r_p$  between EW&AW ranged from 0.20 to 0.76, from 0.89 to 0.96 between EW&YW, from -0.30 to 0.20 between AW & AI and from -0.07 to 0.21 between YW & YI in White Leghorn chickens.

### **Selection effect and generation contrasts**

Estimates of the three generations of selection obtained by generalized least squares using BLUPF90 were superior ( $P < 0.05$ ) for most egg quality traits than the base generation (Table 5). Based on BLUPF90, the contrasts among estimates of these generations were significant ( $P < 0.05$ ). The third generation showed superiority in most egg quality traits compared to the average of the first and second generations. These results agree with Aly *et al.* (2010a), Abou El-Ghar *et al.* (2011), Khalil *et al.* (2013), Shalan *et al.* (2012) and Rayan *et al.* (2013). These superiorities indicted that selection for egg number during the first 90 days of laying in Benha

chickens was associated with improvement in egg component, ESI and EST traits. Shalan *et al.* (2012) found that EW was 42.3, 45.0 and 47.9 g for the base, 1<sup>st</sup> and 2<sup>nd</sup> generations in Baheij selected line and 40.3, 42.9 and 44.0 g for the base, 1<sup>st</sup> and 2<sup>nd</sup> generations in Baheij control line. Younis *et al.* (2014) reported that the averages of EST in Dokki-4 chickens for the base, first and second generations were 0.38, 0.35 and 0.35 mm for the selected line and 0.37, 0.35 and 0.34 mm for the control line, respectively.

#### **The phenotypic and genetic trends across generations:**

The phenotypic trends for egg quality traits across the generations are presented in Figure 1. The initial phenotypic values were 48, 26.8, 14.6 and 6.0 g for EW, AW, YW and SW, respectively. While reached 48, 27.4, 14.6 and 6.0 g in the first generation, 48.5, 27.7, 14.6 and 6.1g in the second generation and 48.9, 28, 28.1, 14.6 and 6.2 g in the third generation, respectively; indicating that egg components were improved by selection. These results fall within the ranges of 42.1 to 52.5 g for EW, from 13.9 to 16.19 g for YW and from 21.6 to 30 for AW as cited in literature (Aly *et al.*, 2010b; Shalan *et al.*, 2012; Alipanah *et al.*, 2013; Taha and Abd El-Ghany, 2013; Khalil *et al.*, 2013; Younis *et al.*, 2014). Phenotypic values of ESI, AI and YI were 76, 7.0 and 42 % in the base generation, respectively (Figure 1). While the phenotypic values were 76.9, 8.0 and 43.0 % in the first generation, 77.4, 8.0 and 44 % in the second generation and 77.9, 9.0 and

45.1% in the third generation, respectively. These results agree with shalan *et al.* (2012), Khalil *et al.* (2013) and Taha and Abd El-Ghany (2013). The initial phenotypic value for haugh unit was 87 in the base generation, while reached 89, 92 and 94 in the first, second and third generation, respectively (Figure 1). These results fall within the ranges showed by Abd El-Latif (2001), Iraqi (2002) and Khalil *et al.* (2013). The phenotypic value for shell thickness initiated at 0.30 mm in the base generation, then increasing slightly till reached 0.31, 0.33 and 0.35 mm in the first, second and third generation, respectively (Figure 1). These trends agree with Samak (2001) and Khalil *et al.* (2013).

The genetic trends across the generations for egg quality traits are presented in Figure 2. The initial BLUP for EW, AW, YW and SW were 0.1, 0.43, -0.19 and 0.09 g in the base generation, while reached 0.4, 0.36, 0.05 and 0.11 g in the first generation, 0.7, 0.30, 0.08 and 0.12 g in the second generation and finally reached 1.0, 0.24, 0.22 and 0.13 g in the third generation, respectively. The BLUP estimates initiated with 0.46, -0.02 and 0.04 % for ESI, AI and EST in the base generation (Figure 2), while reached 0.81, 0.03 and 0.20 % in the first generation, 1.16, 0.07 and 0.35 % in the second generation and 1.52, 0.12 and 0.05 in the third generation, respectively.

#### **Correlated selection responses**

Selection for improving egg number at 90 day increasing EW after three

### **selection, egg quality traits, genetic and phenotypic trends, correlated responses.**

selected generations since the correlated response for EW was 0.44, 0.25 and 0.49 g in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively and 1.18 g as accumulative response (Table 6). The correlated response for AW was 0.11, 0.26 and 0.50 g in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively and 0.87 g with accumulative response, while the YW was 0.38, 0.32 and 0.08 g, in 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively and 0.78 g with accumulative response and the SW was 0.02, 0.27 and 0.02 g in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively with a accumulative response of 0.31g ; these results indicated that the genes that affect the egg number at 90 days are the same genes affecting EW. These results are in agreed with those obtained by Cavero *et al.* (2001) and Shalan *et al.* (2012). Response in HU was -1.11, 5.64 and 0.74 in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation of selection, respectively and 5.27 as accumulative response. The improvement in the HU in the three generations may be due to the increasing in EW and AI values. The correlated response to selection for ESI was -0.42, 2.12 and 0.13 % in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively and 1.83 % with accumulative response. Response in AI was 0.47, -1.55 and 2.45% in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation, respectively with accumulative response of 1.37 %). These results may be due to that increasing EW after three generations of selection was associated with improvement in egg components. Yolk index was 0.24, 0.75 and 0.96 % in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generation,

respectively with accumulative response of 1.95%. This result in accordance with Younis *et al.* (2014). The responses in EST were 0.04, 0.01 and 0.01mm in the 1<sup>st</sup>, 2<sup>nd</sup> and 3<sup>rd</sup> generations, respectively and of 0.06 as a cumulative response. Similar trend was observed by Younis *et al.* (2014) and Shalan *et al.* (2012).

### **CONCLUSIONS**

- 1) The high estimates of BLUP for egg quality traits in the control and selected generations indicate that improvement of egg quality traits in Benha chickens could be achieved through more generations of selection for egg number during the 1<sup>st</sup> 90 days of laying
- 2) Superiorities in the selected generations indicated that selection for egg number during the first 90 days of laying in Benha chickens was associated with an improvement in egg quality traits
- 3) The genetic trends were slightly positive for most traits indicating that the selection program was performed correctly.

**Table (1):** Numbers (No.) sires, dams, pullets and eggs used.

Generation	No. of sires	No. of dams	No. of Pullets	No. of eggs used
Base (control) population	17	113	180	603
1 <sup>st</sup> generation of selection	18	119	180	1476
2 <sup>nd</sup> generation of selection	17	117	180	897
3 <sup>rd</sup> generation of selection	17	135	216	1266
TOTAL	69	484	756	4242

**Table (2):** Actual means, standard deviations (SD), and percentages of variation (V%) for egg quality traits in Benha chickens.

Trait	Symbol	Mean	SD	V%
Egg weight (g)	EW	48.3	4.7	9.8
Albumen weight (g)	AW	27.6	3.4	12.5
Yolk weight (g)	YW	14.6	2.0	13.9
Shell weight (g)	SW	6.1	0.85	13.9
Haugh unit	HU	90.6	26.0	28.6
Egg shape index (%)	ESI	78.4	0.17	0.21
Albumen index (%)	AI	8.0	0.02	0.24
Yolk index (%)	YI	44.0	0.05	0.11
Egg shell thickness (mm)	EST	0.33	0.04	0.12

**Table (3):** Estimates of additive genetic ( $\sigma^2_a$ ), permanent environment ( $\sigma^2_{PE}$ ), uncontrolled environment ( $\sigma^2_E$ ) and phenotypic ( $\sigma^2_p$ ) variances along with heritability estimates ( $h^2 \pm SE$ ) for egg quality traits in Benha chickens.

Trait	$\sigma^2_a$	$\sigma^2_{PE}$	$\sigma^2_E$	$\sigma^2_p$	$h^2 \pm SE$
EW (g)	16.1	8.4	9.5	34	0.47±0.02
AW (g)	11	6.4	9.7	27.1	0.40±0.02
YW (g)	1.1	0.97	1.0	3.0	0.36±0.02
SW (g)	0.75	2.1	1.6	4.4	0.17±0.02
HU	4.5	25.1	44.3	74	0.06±0.02
ESI (%)	1.7	2.2	2.9	6.9	0.25±0.01
AI (%)	0.6	1.8	2.7	5.1	0.11±0.01
YI (%)	0.07	0.5	0.84	1.4	0.05±0.01
EST (mm)	2.1	2.8	4.2	9.1	0.23±0.01



**Table (4):** Estimates of genetic ( $r_G$ ), and phenotypic ( $r_P$ ) correlations among some egg quality traits in Benha chickens.

Traits correlated	$r_G$	$r_P$
EW & AW	0.99±0.01	0.85±0.17
EW & YW	0.57±0.11	0.63±0.14
EW & SW	0.69±0.13	0.54±0.16
AW&AI	0.59±0.10	0.02±0.08
YW&YI	0.61±0.11	-0.26±0.13

635

**Table (5):** generalized Least-square means for egg quality traits in different generations along with all possible generation contrasts and their standard errors ( $\pm$  SE) in Benha chickens.

Generations	EW (g)	AW (g)	SW (g)	HU	ESI (%)	STH (mm)
Base generation ( $G_0$ )	47.4±0.19 <sup>c</sup>	27.3±0.14 <sup>b</sup>	6.0±0.03 <sup>b</sup>	88.5±1.05 <sup>b</sup>	76.7±0.007 <sup>b</sup>	0.27±0.01 <sup>b</sup>
1 <sup>st</sup> generation ( $G_1$ )	48.1±0.12 <sup>b</sup>	27.3±0.09 <sup>b</sup>	5.9±0.02 <sup>b</sup>	87.3±0.64 <sup>b</sup>	76.3±0.005 <sup>b</sup>	0.32±0.01 <sup>c</sup>
2 <sup>nd</sup> generation ( $G_2$ )	48.3±0.15 <sup>b</sup>	27.5±0.11 <sup>b</sup>	6.2±0.02 <sup>a</sup>	92.9±0.86 <sup>a</sup>	78.9±0.006 <sup>a</sup>	0.33±0.01 <sup>a</sup>
3 <sup>rd</sup> generation ( $G_3$ )	49.1±0.13 <sup>a</sup>	28.1±0.10 <sup>a</sup>	6.2±0.02 <sup>a</sup>	93.8±0.72 <sup>a</sup>	79.1±0.005 <sup>a</sup>	0.34±0.01 <sup>a</sup>
$G_0$ VS $G_1$	-0.7±0.22*	0.0±0.0	0.1±0.04	1.2±1.40	0.4±0.009	0.05±0.01-
$G_0$ VS $G_2$	-0.9±0.24*	-0.2±0.18	-0.2±0.04*	-4.4±1.36*	-2.2±0.009*	-0.0±0.03*
$G_0$ VS $G_3$	-1.7±0.23*	-0.8±0.17*	-0.2±0.04*	-5.3±1.27*	-2.4±0.009*	-0.01±0.003*
$G_1$ VS $G_2$	-0.2±0.02	-0.3±0.14	-0.3±0.03*	-5.6±1.09*	-2.6±0.008*	-0.01±0.002*
$G_1$ VS $G_3$	-1.0±0.18*	-0.8±0.13*	-0.3±0.03*	-6.5±0.99*	-2.8±0.007*	-0.02±0.002
$G_2$ VS $G_3$	-0.8±0.20*	-0.6±0.15*	0.0±0.0	-0.9±1.12	-0.2±0.008	-0.01±0.002

a, b, c, d Means with the same letters within each trait are not significantly different ( $P < 0.05$ ).


\* Generation effect significantly different from 0,  $\alpha = 0.05$ .

**Table (6):** Realized correlated response for unselected egg components traits in Benha chickens through three generations of selection.


Traits	Correlated selection response as a result of selection for egg number during the 1 <sup>st</sup> 90 day of laying (EN90D)			
	1 <sup>st</sup> generation	2 <sup>nd</sup> generation	3 <sup>rd</sup> generation	cumulative
EW (g)	0.44	0.25	0.49	1.18
AW (g)	0.11	0.26	0.50	0.87
YW (g)	0.38	0.32	0.08	0.78
SW (g)	0.02	0.27	0.02	0.31
HU	-1.11	5.64	0.74	5.27
ESI (%)	-0.42	2.12	0.13	1.83
AI (%)	0.47	-1.55	2.45	1.37
YI (%)	0.24	0.75	0.96	1.95
EST (mm)	0.04	0.01	0.01	0.06

**selection, egg quality traits, genetic and phenotypic trends, correlated responses.**

**Figure (1):** The phenotypic trend for egg quality traits in Benha chickens across the selected generations.



**Figure (2):** Genetic trend for egg quality traits in Benha chickens across the selected generations.



REFERENCES

- Abdellatif, M. A. 2001.** Inheritance of certain plasma constituents and their association with some economic traits in Dandarawi and Golden Montazah hens. M.Sc. Thesis, Fac. Of Agric., Cairo Univ., Egypt.
- Abou El-Ghar, R. Sh.; Aly, O. M. and Ghanem, H. H. 2011.** Test of epistasis among triple crosses of Matrouh with different strains of chickens. Egyptian Poultry Science, 31: 29-38.
- Alipanah, M.; Deljo, J.; Rokouie, M. and Mohammadnia, R. 2013.** Heritabilities and genetic and phenotypic correlations of egg quality traits in khazak layers. Trakia Journal of Sciences, 2:175-180.
- Aly, O. M. ; Ghanem, Hanan H.; Afifi, Yousria K.; Abou El-Ella, Nazla Y. and Balat, Magda M. 2010a.** Selection for improving egg production in Mandarah chickens. 4-Direct and correlated response for some economic traits for four generations of selection. Egyptian Poultry Science, 30: 137-156.
- Aly, O. M.; Mosaad, Nema A.; Abou El-Ella, Nazla Y. and Afifi, Yousria K. 2010b.** Improving the productivity and productivity of Bahij chickens through crossing. Effect of upgrading on: B-Egg production, egg quality and hatchability traits. Egyptian Poultry Science, 30: 1151- 1170.
- Ashour, A. F.; Badwi, Y. K. and Abd El-Karim, Ragaa E. 2015.** Effect of selection for body weight on egg production, egg quality, fertility and hatchability traits in El-Salam chickens strain in Egypt. J. Animal and poultry Prod., Mansoura Univ., 6 (12): 781-796.
- Cavero, D.M.; Schmutz, Ieken. K and Presinger, R 2001.** Improving hatchability in white egg layer strains through breeding. Lohmann information, 46 (1): 44-54.
- Dasari, S.; Manthani, G.P.; Mallam, M. and Rudra, N.C. 2013.** Genetic analysis of egg quality traits in White Leghorn chicken. Veterinary world, 6(5) :263-266.
- Dunn, I.C.; Bain, M.; Edmond, A.; Wilson, P.W.; Joseph, N.; Solomon, S.; De Ketelaere, B.; De Baerdemaeker, J.; Schmutz, M.; Preisinger, R. and Waddington, D. 2005.** Heritability and genetic correlation of measurements derived from acoustic resonance frequency analysis; a novel method of determining eggshell quality in domestic hens. British poultry science, 46(3) : 280-286.
- El- Garhy, O. H. 2004.** Effect of some biotechnological methodology on some productive and reproductive traits in fowl. M.Sc. Thesis, Fac. Of Agric. Moshtohor, ZagazigUniv, Benha.Branch, Egypt.
- El-Attrouny, M. M. 2011.** Estimation and evaluation of genetic and non-genetic parameters for some productive traits in chickens. M. Sc. Thesis, Faculty of Agriculture at Moshtohor, Benha University, Egypt.
- El-Attrouny, M.M. and Iraqi, M.M. 2021.** Influence of selection for egg production on egg quality traits in Japanese quail. South African Journal of Animal Science, 51(1):128-137
- El-Attrouny, M.M.; Iraqi, M.M.; Khalil, M.H. and El-Moghazy, G.M. 2017.** Genetic and phenotypic evaluation of growth traits in selection experiment performed in synthesized Benha chickens. Annals of Agric. Sci., Moshtohor, 51(1): 33-42.

- El-Attrouny, M.M.; Khalil, M.H.; Iraqi, M.M. and El-Moghazy, G.M. 2019.** Genetic and Phenotypic evaluation of egg production traits in selection experiment performed on Benha Chickens. *Egyptian Poultry Science Journal*, 39(2):459-477.
- Falconer, D. S. and Mackay, F. C. 1996.** Introduction to Quantitative Genetics. "4<sup>th</sup> Edition" Longman Group Ltd, England.
- Flock, D.K., Schmutz, M. and Preisinger, R. 2007.** Optimization of egg quality from the breeders, point of view. *Zuchtungskunde*, 79(4):309-319.
- Hanafi, M. S. and El-Labban, A. F. 1990.** Genetic studies on egg contents and shell characteristics in Dokki-4 chickens. *Agricultural Research Review*, 68 (4): 669-681.
- Harms, R.H.; Douglas, C.R. and Sloan, D.R. 1996.** Midnight feeding of commercial laying hens can improve eggshell quality. *Journal of Applied Poultry Research*, 5(1):1-5.
- Hunton, P. 1995.** Understanding the architecture of the egg shell. *World's Poultry Science Journal*, 51(2):141-147.
- Icken, W.; Schmutz, M. and Preisinger, R. 2006.** Dynamic stiffness measurements with the "crack detector": a new method to improve egg shell strength.
- Iraqi, M. M. 2002.** Genetic evaluation for egg quality traits in crossbreeding experiment involving Mandarah and Matrouh chickens using Animal Model. *Egyptian Poultry Science*, 22. (3): 711-726.
- Iraqi, M. M.; Khalil, M. H. and El-Attrouny, M. M. 2013.** Estimation of crossbreeding components for growth traits in crossing Golden Montazah with White Leghorn chickens. VI<sup>th</sup> International Conference: Balnimalcon 2013 Tekirdag/Turkiye, October 3-5: 494-504.
- Kennedy, B. W. 1989.** Animal model BLUP - Crasmus intensive graduate Course. University of Guelph, Dublin.
- Khalil, M. H.; Iraqi, M. M. and El-Attrouny, M. M. 2013.** Effects on egg quality traits of crossing Egyptian Golden Montazah with White Leghorn chickens. *Livestock Research for Rural*, 25 (6): 2013.
- Khawaja, T.; Khanb, S. H.; Mukhtara, N.; Ullaha, N. and Parveenb, A. 2013.** Production performance, egg quality and biochemical parameters of Fayoumi, Rhode Island Red and their reciprocal crossbred chickens. *Journal of Applied Animal Research*, 41, 2: 208- 217.
- Misztal, I.; Lourenco, D.; Aguilar, I.; Legarra, A. and Vitezica, Z. 2014.** Manual for BLUPF90 family of programs. University of Georgia, Athens, USA. First released May 12, 2014 Last Edit June 18, 2015.
- NRC, 1994.** National Research Council Nutrient Requirement of Poultry. Ninth Revised Ed. National Academy Press Washington DC, USA.
- Olawumi, S. O. and Jobin. A. 2013.** Phenotypic correlation between live body weight and carcass traits in Arbor Acre breed of broiler chickens. *International Journal of Science and Nature*, 4: 145-149.
- Olawumi, S. O. and Ogunlade, J. T. 2008.** Phenotypic correlation between some external and internal egg quality traits in the exotic Isa Brown layer breeders. *Asian Journal of poultry Science* 2 (1): 30-35.
- Preisinger, R. and Flock, D. K. 2000.** Genetic changes in layer breeding: historical trends and future

**selection, egg quality traits, genetic and phenotypic trends, correlated responses.**

---

- prospects. BSAP Occasional Publication, 27:20-28.
- Rath, P. K.; Mishra, P. K.; Mallick, B. K. and Behura, N. C. 2015.** Evaluation of different egg quality traits and interpretation of their mode of inheritance in White Leghorns. *Veterinary World*, ISSN: 2231-0916.
- Rayan, G. N.; Mahrous, M. Y.; Galal, A. and El-Attar, A. H. 2013.** Study of some productive performance and egg quality traits in two commercial layer strains. *Egyptian Poultry Science*, 33 (2): 357-369.
- Samak, H. R. 2001.** Evaluating and improving the productive performance in some local chickens. Ph.D. Thesis, Fac. of Agric., Zagazig. Univ. Benha branch.
- Shalan, Hedaia.M.; El-Sayed, Nadia A. and Rizk, R. E. 2012.** Correlated response in shell and albumin quality in local chicken strain selected for increasing egg mass. *Egyptian Poultry Science*, 32 (11): 593-605
- Taha, A. E. and Abd El-Ghany, F. A. 2013.** Improving Production Traits for El-Salam and Mandarah Chicken Strains by Crossing. I-Estimation of crossbreeding effects on egg production and egg quality traits. *World Academy of Science Engineering and Technology, International Journal of Biological Science and Engineering* 7 (10).
- Toussant, M. J. and Latshaw, J. D. 1999.** Ovomucin contents and composition in chicken eggs with different interior quality. *J. Sci. Food and Agri.*, 79:1666-1670.
- Udoh, U. H.; Okon, B. and Udoh, A.P. 2012.** Egg quality characteristics, Phenotypic Correlations and Prediction of egg weight in three (Naked Neck, Frizzled Feather and normal Feathered) Nigerian Local Chickens. *International Journal of Poultry Science*, 11 (11): 696-699.
- Younis, H. H.; Abd El-Ghany, F. A. and Awadein, Nasra B. 2014.** Genetic improvement of egg production traits in Dokki-4 strain. 1- Correlated response, heritability, genetic and phenotypic correlations for egg production and egg quality traits. *Egyptian Poultry Science*, 34:345-362.
- Zhang, L.C.; Ning, Z.H.; XU, G.Y.; Choum, Z. and Yong, N. 2005.** Heritability and genetic and phenotypic correlation of egg quality traits in brown egg dwarf layers. *Poult. Sci.* 84: 1209-1213.

## الاستجابة المرتبطة لصفات جوده البيض بعد ثلاث اجيال من الانتخاب لعدد البيض فى دجاج بنها

محمود مصطفى الاطرونى<sup>١</sup>، محمود مغربى عراقى<sup>١</sup>، ماهر حسب النبى خليل<sup>١</sup>، جيهان محمد المغازى<sup>٢</sup>

<sup>١</sup>قسم الإنتاج الحيوانى كلية الزراعة بمشهر – جامعة بنها - مصر  
<sup>٢</sup>المركز الإقليمي للأغذية والاعلاف – مركز البحوث الزراعية – مصر

هدفت الدراسة الحالية علي دراسة تأثير الانتخاب لزيادة عدد البيض خلال ال ٩٠ يوم الاولى من الوضع علي صفات جودة البيض خلال أربعة اجيال فى دجاج بنها. تم إنتخاب ١٨ ديك و ١٨٠ دجاجة طبقاً لقيم أحسن متنبئ خطى غير متحيز BLUP لصفة عدد البيض عند اول ٩٠ يوم من الإنتاج من جيل الأساس (الكنترول). تم جمع عدد 4242 بيضة ناتجة من ٧٥٦ دجاجة ناتجة من ٦٩ ذكر و ٤٨٤ أم للأربع أجيال (جيل أساسى وثلاث أجيال إنتخابية) لتقدير المكافئ الوراثى – الارتباط الوراثى والمظهرى وقيم أحسن متنبئ خطى غير متحيز BLUP لصفات جودة البيضة (وزن البيضة، وزن البياض، وزن الصفار، وزن القشرة، وحدات هو، دليل شكل البيضة، دليل البياض، دليل الصفار وسمك القشرة). تم تقدير أوتوضيح أثر الانتخاب والإستجابة المرتبطة عبر الاجيال والإتجاهات الوراثية والمظهرية لصفات جودة البيض فى دجاج بنها بإستخدام نموذج الحيوان المعروف باسم BLUPF90 Program. كانت تقديرات المكافئ الوراثى قيماً متوسطة او منخفضة لصفات جودة البيض حيث كانت بين ٠.٠٥ الي ٠.٤٧. ايضاً أظهرت الارتباطات الوراثية قيم موجبة بين صفة وزن البيض وكلاً من وزن البياض، وزن الصفار ووزن القشرة حيث كانت ٠.٩٩، ٠.٥٧، ٠.٦٩. على التوالى.. أظهرت الأجيال الثلاثة الإنتخابية تفوقاً لمعظم صفات جودة البيض عن جيل الأساس وكانت الفروق معنوية عند مستوى معنوية ( $P<0.05$ ). كما أظهرت التضادات المستقلة Contrasts بين الأجيال المختلفة اختلافاً معنوياً. أظهر الإتجاهات المظهرية والوراثية إزدياداً لمعظم صفات جودة البيض. بينما انخفض لصفة وزن البياض. فى القيم المظهرية. أظهرت الإستجابة الإنتخابية المحققة والمصاحبة بعد ثلاث أجيال إنتخابية زيادة تراكمية لكلاً من الصفات التالية: وزن البيضة (١.١٨ جرام)، وزن القشرة (٠.٣١ جرام)، وحدات هو (٥.٢٧) وسمك القشرة (٠.٠٦ ملليمتر) على التوالى. الخلاصة، تشير هذه النتائج إلى أن تفوق الأجيال المنتخبة يشير إلى أن انتخاب عدد البيض خلال التسعين يوماً الأولى من الوضع فى دجاج بنها كان مرتبطاً بتحسين صفات جودة البيض.