

Proceedings of the



4-7 July 2000 – Valencia Spain

These proceedings were printed as a special issue of WORLD RABBIT SCIENCE, the journal of the World Rabbit Science Association, Volume 8, supplement 1

ISSN reference of this on line version is 2308-1910

(ISSN for all the on-line versions of the proceedings of the successive World Rabbit Congresses)

KHALIL M.H., AFIFI E.A.

**HETEROSIS, MATERNAL AND DIRECT ADDITIVE EFFECTS
FOR LITTER PERFORMANCE AND POSTWEANING GROWTH
IN GABALI RABBITS AND THEIR F₁ CROSSES
WITH NEW ZEALAND WHITE**

Volume A, pages 431-437

HETEROSIS, MATERNAL AND DIRECT ADDITIVE EFFECTS FOR LITTER PERFORMANCE AND POSTWEANING GROWTH IN GABALI RABBITS AND THEIR F₁ CROSSES WITH NEW ZEALAND WHITE

KHALIL MH.*, AFIFI EA.**

*Department of Animal Production and Breeding, College of Agriculture and Veterinary Medicine, King Saud University, Buriedah, P.O. Box 1482, Saudi Arabia

** Department of Animal Production, Faculty of Agriculture at Moshtohor, Zagazig University, Egypt.

ABSTRACT

: A crossbreeding experiment was carried out involving Gabali (**G**) and New Zealand White (**NZW**) rabbits to estimate direct heterosis (**H^I**) and direct (**G^I**) and maternal (**G^M**) additive effects on some litter traits and postweaning growth. Data on litter traits of 314 litters (litter size and weight at birth and weaning, milk yield at 21 days and total milk yield) and body weight (at 4, 6, 8, 10 and 12 weeks of age) of 1300 weaned rabbits were analyzed using a linear mixed model. **G^I** did not significantly affect most litter traits and postweaning growth. **G**-sired litters had similar direct additive effects compared to **NZW**-sired litters and consequently **G** bucks could be used as sires in crossbreeding stratification systems under hot climatic conditions. Crossbred litters (or rabbits) obtained from mating **G** bucks with **NZW** does were generally associated with slight superiority compared to those litters (or rabbits) obtained from the reverse mating. The estimates of **G^M** for litter size and weight at birth were significantly in favour of **G**-rabbits, while breed maternity for litter traits measured after kindling was significantly in favour of **NZW** breed. After weaning, growth traits were not significantly affected by **G^M**. Crossing of **G** rabbits with **NZW** was associated with significant positive estimates of **H^I** for litter size and weight at birth and weaning. Slight negative estimates of **H^I** were observed for milk yield at 21 days and total milk yield. However, insignificant negative estimates of **H^I** were recorded for postweaning growth traits.

INTRODUCTION

Gabali rabbits raised under the Egyptian desert conditions (especially in Sinai) are characterized by a large litter size of 8-12 young and heavy body weight of 3.5-4.5 kg (GALAL and KHALIL, 1994). Crossbreeding between standard breeds and Gabali rabbits raised under the desert conditions is not widely carried out. To date, publications concerning crossbreeding of Gabali rabbits with standard breeds (e.g. New Zealand White) in Egypt are not available. Direct and maternal heterosis, maternal and direct additive effects from crossbreeding experiments including Gabali rabbits were expected to be important especially for post-weaning growth performance (KHALIL, 1996). On the other hand, the New Zealand White breed was found to exhibit an outstanding maternal ability as related to behavior, fecundity and lactation (LUKEFAHR et al, 1983ab; OZIMBA and LUKAFAHR, 1991). Results of most crossbreeding experiments carried out in Egypt reported that crossing does of New Zealand White breed with bucks of local breeds were generally associated with considerable heterotic effects on most litter and growth traits (OUDAH, 1990; EL-DESOKI, 1991; AFIFI et al, 1994; KHALIL et al, 1995). Therefore, this study was conducted to estimate direct (**G^I**) and maternal (**G^M**) additive effects and direct heterosis (**H^I**) for some litter traits and postweaning body weights in a crossbreeding experiment involving New Zealand White and Gabali rabbits.

MATERIAL AND METHODS

A crossbreeding experiment was carried out in the Experimental Rabbitry at Moshtohor, Zagazig University, Egypt (about 27 Km to the north of Cairo) during one production year started in November 1994.

Rabbits used in this study represent one desert Egyptian breed (Gabali, **G**) and one exotic breed (New Zealand White, **NZW**). Each buck was allowed to sire all his litters from the same does. The breeding plan permitted the simultaneous production of **G**, **NZW**, **G x NZW** and **NZW x G** litters in each parity. Matings in this experiment started in November 1994 and stopped in April 1995. Distribution of breeding does and bucks and number of litters and bunnies born and weaned of the four different genetic groups are presented in Table 1.

Rabbits were raised in a semi-closed rabbitry. Breeding does and bucks were housed separately in individual wired-cages.. Does were mated from the same assigned bucks 10 days after each kindling. Young rabbits were weaned at four weeks, ear tagged, sexed and transferred to standard progeny wire cages equipped with feeding hoppers and drinking nipples. Feeding practices in the rabbitry were described by KHALIL (1994).

Data of litter traits included litter size at birth (**LSB**) and weaning (**LSW**), litter weight at birth (**LWB**) and weaning (**LWW**), milk yield at 21 days (**M21**) and total milk yield (**TMY**), while data of postweaning growth included body weights at 4, 6, 8, 10 and 12 weeks of age (**W4**, **W6**, **W8**, **W10** and **W12**, respectively). Data of litter traits on 314 litters (Table 1) were analyzed using the following mixed model (HARVEY, 1990):

$$Y_{ijklm} = \mu\mu + G_i + D_{ij} + A_k + P_l + e_{ijklm} \quad (\text{Model 1})$$

Where Y_{ijklm} is the observation on the $ijklm^{\text{th}}$ litter trait (**LSB**, **LSW**, **LWB**, **LWW**, **M21** and **TMY**); $\mu\mu$ is the overall mean; G_i is the fixed effect of i^{th} breed group; D_{ij} is the random effect of j^{th} doe nested within the i^{th} breed group, A_k is the fixed effect of the k^{th} season of kindling ($k=1,2$); P_l is the fixed effect of l^{th} parity ($l=1,\dots,5$); and e_{ijklm} is the random deviation particular to the m^{th} litter, **NID** ($0, \sigma\sigma^2_e$).

Table 1. Number of bucks, does, litters and bunnies distributed in the four breed groups of the study

Genetic group ⁺	Bucks	Does	Litters weaned	Bunnies born	Bunnies weaned
NZWxNZW	15	50	160	1173	850
GxG	8	26	56	348	280
NZWxG	8	25	60	524	360
GxNZW	14	37	98	698	580
Total	55	138	374	2743	2070

⁺Breed of buck is listed before breed of doe.

Data on 2070 weaned (Table 1) rabbits for postweaning growth traits were analyzed using the following sire model (HARVEY, 1990):

$$Y_{ijklmno} = \mu\mu + G_i + S_{ij} + A_k + P_l + B_m + C_n + (GB)_{im} + (AP)_{kl} + (AB)_{km} + (AC)_{kn} + (PB)_{lm} + (BC)_{mn} + e_{ijklmno} \quad (\text{Model 2})$$

Where $Y_{ijklmno}$ is the observation on the $ijklmno^{\text{th}}$ weaned rabbit of the postweaning growth trait (**W4**, **W6**, **W8**, **W10** and **W12**); $\mu\mu$ is the overall mean, G_i is the fixed

effect of the i^{th} breed group, S_{ij} is the random effect of the j^{th} sire nested within the i^{th} breed group (taking the relationship coefficient inverse matrix A^{-1} into consideration), A_k is the fixed effect of the k^{th} season of birth ($k=1,2$); P_l is the fixed effect of the l^{th} parity ($l=1,\dots,5$); B_m is the fixed effect of the m^{th} sex; C_n is the fixed effect of the n^{th} teats number of doe ($n= 5,6,\dots 10$); and $e_{ijklmno}$ is the random deviation particular to the o^{th} weaned rabbit, NID ($\mathbf{0}, \sigma\sigma_e^2$) along with all possible interactions of GB_{im} , AP_{kl} , AB_{km} , AC_{kn} , PB_{lm} and BC_{mn} . Breed group was tested against sire within breed group, while other fixed effects were tested against the remainder.

Crossbreeding effects of direct additive (G^I), maternal additive (G^M) and direct heterosis (H^I) for different litter traits and body weights were estimated according to Dickerson theory (DICKERSON, 1992). Such genetic model permits to derive a selected set of linear contrasts and therefore G^I , G^M and H^I were estimated as:

Direct additive effect:

$$(G^I_{NZW} - G^I_G) = \{[(NZW \times NZW) + (NZW \times G)] - [(G \times G) + (G \times NZW)]\}$$

Maternal additive effect:

$$(G^M_{NZW} - G^M_G) = [(G \times NZW) - (NZW \times G)]$$

Direct heterotic effect (units):

$$H^I \text{ in units} = [(NZW \times G + G \times NZW) - (NZW \times NZW + G \times G)]/2$$

$$H^I (\%) = [(NZW \times G + G \times NZW) - (NZW \times NZW + G \times G)] / [(NZW \times NZW + G \times G)(100)]$$

Where G^I and G^M represent direct additive and maternal additive effects, respectively, of the subscripted genetic group. Each single degree of freedom contrast was tested for significance with the Student's t-test.

RESULTS AND DISCUSSION

Direct additive effect (G^I)

The linear contrasts of G^I for most litter and postweaning growth traits were insignificant (Tables 2&3). Such limited differences in G^I between the two breeds lead to state that G could be used as a buck-breed in crossbreeding programmes.

NZW had higher estimates of G^I than G -for litter weights at birth ($P < 0.001$ and at weaning ($P < 0.10$). High estimates of G^I for litter traits lead to indicate that NZW breed could be used as a terminal sire breed for litter traits measured at kindling. In France, an experiment (1970) showed that Californian-sired litters had higher estimates of G^I for pre-weaning litter traits than that of NZW -sired litters (ROUVIER and BRUN, 1990). A reverse trend was observed in an experiment performed 20 years later (BRUN, 1993). Also, the American study by LUKEFAHR et al (1983a) showed that estimates of G^I for pre-weaning litter traits (**LSB**, **LWB**, **LSW** and **LWW**) were mostly in favour of Californian vs NZW . They added that direct Flemish Giant effects on pre-weaning litters were positive and high compared with NZW .

For post-weaning growth traits the G^I effects were only significant for the weight at 12 weeks, being the value of G breed higher than the one of the NZW (Table 3). Such superiority of G -sired rabbits in G^I may be an encouraging factor for the rabbit breeder in hot climate countries to use their native breeds in any crossbreeding stratification system. Estimates of G^I presented in Table 5 indicate also that G^I at later age (**W12**) was significantly in favour of G breed. At later ages, MASOERO et al (1985) evidenced such significant G^I in NZW , Californian, Burgundy Fawn, Flemish Giant, Argenté de Champagne and Blue Vienna and their crosses.

Table 2. Genetic group means (+SE) and estimates of direct additive effect (G^I), maternal additive effect (G^M) and direct heterosis (H^I) of litter traits⁺

Item	LSB	LWB	LSW	LWW	M21	TMY
Genetic group⁺⁺:						
NZWxNZW	6.75±0.22	403±13. 7	4.58±0.24	2849±143	2320±92	3482±13 6
GxG	5.91±0.81	305±50. 2	3.49±0.89	2206±524	2235±33 1	3383±49 7
NZWxG	8.32±0.82	444±50. 4	4.63±0.89	2746±526	1963±32 6	3111±49 7
GxNZW	6.65±0.49	389±30. 5	4.97±0.54	3099±318	2384±19 6	3512±30 0
Significance	*	NS	NS	NS	NS	NS
Direct additive effect:	2.52 ^{***}	152 ^{***}	0.8 ^{NS}	291 ^{NS}	-336 ^{NS}	-598 ^{NS}
Maternal additive effect:	-1.7 ^{***}	-55 [*]	0.3 ^{NS}	353 [*]	421 [*]	401 [*]
Heterosis						
H ^I	1.2 ^{***}	62 ^{**}	0.8 ^{**}	395 [*]	-104 ^{NS}	-122 ^{NS}
H ^I (%)	18.3	17.5	18.8	15.6	-4.6	-3.5

⁺⁺Buck-breed listed first NS= Non-significant (P>0.05); *=P<0.05; **=P<0.01; ***=P<0.001.

Table 3. Genetic group means (+SE) and estimates of direct additive effect (G^I), maternal additive effect (G^M) and direct heterosis (H^I) for post-weaning body traits⁺

Item	W4	W6	W8	W10	W12
Genetic group⁺⁺:					
NZWxNZW	591±22	807±23	1065±30	1378±35	1711±36
GxG	587±56	796±62	1084±83	1405±95	1812±87
NZWxG	572±99	820±83	1073±105	1411±114	1718±103
GxNZW	584±51	798±52	1063±66	1364±76	1726±72
Significance	NS	NS	NS	NS	NS
Direct additive effect:	-7.9 ^{NS}	32.7 ^{NS}	-8.6 ^{NS}	19.2 ^{NS}	-108.4 [*]
Maternal additive effect:	12.2 ^{NS}	-21.8 ^{NS}	-10.1 ^{NS}	-47.1 ^{NS}	7.8 ^{NS}
Heterosis :					
H ^I	-10.4 ^{NS}	-7.5 ^{NS}	-6.5 ^{NS}	-3.7 ^{NS}	-39 ^{NS}
H ^I (%)	-1.8	-0.9	-0.6	-0.3	-2.2

⁺⁺ Sire-breed listed first NS= Non-significant (P>0.05); *= P<0.05.

Maternal additive effect (G^M)

Estimates of G^M for litter sizes and weights at birth were significant and they were mainly in favour of G breed (Table 2). After kindling, G^M was significantly in favour of NZW breed for M21, TMY and LWW. This superiority of NZW does is attributable to favorable maternal abilities and an increased levels of milk production compared to G does. Crossbreeding experiments carried out in Egypt (AFIFI and KHALIL, 1989; OUDAH, 1990; KHALIL et al, 1995) indicated that estimates of G^M for pre-weaning

litter traits were insignificant. However, most of the Egyptian findings reported a general trend indicating that litters mothered (*direct plus maternal additive effects*) by exotic breeds (e.g. NZW, Californian, Chinchilla... etc.) recorded better performance than litters mothered by native breeds (e.g. Giza White and Baladi rabbits). This evidenced the superiority of exotic breeds in their maternity (in terms of milk production, maternal behavior and care for young). For most pre-weaning litter traits, maternal superiority of NZW breed compared with other standard breeds has been demonstrated in the American studies (e.g. LUKEFAHR et al, 1983ab; OZIMBA and LUKAFAHR, 1991) and in the European studies (e.g. PARTRIDGE et al, 1981; MASOERO et al, 1985; ROUVIER and BRUN, 1990), i.e. using NZW as a dam breed produced high performances in litter size and weight compared to other dam breeds.

Maternal additive effects on all postweaning body weights were insignificant (Table 3). In Egypt, AFIFI et al (1994) found that postweaning growth of rabbits mothered by NZW breed are nearly similar to those rabbits mothered by Baladi Red breed.

Direct heterosis (H^I)

Estimates of H^I (calculated in actual units and as percentages) for different traits are given in Tables 2&3. The estimates indicated also that crossing between NZW and G rabbits was usually associated with an existence of heterotic effects on litter size and weight measured at kindling and at weaning. Estimates of H^I were significant for LSB, LWB, LSW and LWW. Different crossbreeding experiments carried out in Egypt (AFIFI and EMARA, 1984; AFIFI and KHALIL, 1989; KHALIL et al, 1995) revealed that heterotic effects for litter size and litter weight were evidenced.

Estimates of H^I for LSB, LWB, LSW and LWW were positive and ranged from 15.6 to 18.8%. Estimates for milk production (M21 and TMY) and postweaning body weights (W4, W6, W8, W10 and W12) were negative and low.

Conclusions

(1) Since post-kindling litter performances in New Zealand White and Gabali rabbits were not significantly different in their breed performance, one may use either of the two breeds as sires. For rabbits industry, Gabali bucks could be used in terminal crossbreeding system especially in areas of hot climate.

(2) Single cross resulted from mating Gabali sires with New Zealand White dams is recommended and producers and processors could attained economic benefits of commercial production through using of this simple cross.

REFERENCES

- AFIFI EA., EMARA ME. 1984: Litter weight in local Egyptian and exotic breeds of rabbits and their crosses. *Proceeding 3rd World Rabbit Congress, 4-8 April 1984, Rome, Italy, Vol. QGI:126-135.*
- AFIFI EA., KHALIL MH. 1989: Observations on purebred and crossbred litters of Giza White and Grey Giant Flander rabbits in Egypt. *Journal of Applied Rabbit Research, 12: 273-277.*
- AFIFI EA., KHALIL, MH., KHADR AF., YOUSSEF YMK. 1994: Heterosis, maternal and direct effects for postweaning growth traits and carcass performance in rabbit crosses. *Journal of Animal Breeding and Genetics, 111: 138-147.*

- BRUN JM1993: Parameters du croisement entre 3 souches de lapin et analyse de la reponse a une selection sur la taille de portee: caracteres des portees a la naissance et au sevrabr. *Genet.Sel. Evol.*, **25**: 459-474.
- DICKERSON GE. 1992: Manual for evaluation of breeds and crosses of domestic animals. *Food and Agriculture Organization of the United Nations*, Rome, pp 47.
- EL-DESOKI AEM. 1991: Study of the effect of some genetic and environmental factors affecting meat yield from some foreign and local breeds of rabbits and their crosses. *M.Sc. Thesis, Faculty of Agriculture, Mansoura University, Egypt*.
- GALAL ESE., KHALIL MH. 1994: Development of rabbit industry in Egypt. *Options Mediterraneennes*, **8**: 43-56.
- HARVEY WR. 1990: User's Guide for LSMLMW. Mixed model least squares and maximum likelihood computer program. PC-Version 2, *Ohio State University, Columbus, USA (Mimeograph)*.
- KHALIL MH. 1994: Lactational performance of Giza White rabbits and its relation with preweaning litter traits. *Animal Production*, **59**: 141-145.
- KHALIL MH. 1996: Breeds of rabbits in desert regions. *Symposium on Development of rabbits in Siani. Siani, 24-27 September 1996, Egypt (In Arabic)*.
- KHALIL MH., AFIFI EA., YOUSSEF YMK., KHADR, AF. 1995. Heterosis, maternal and direct genetic effects for litter performance and reproductive intervals in rabbit crosses. *World Rabbit Science*, **3(3)**: 99-105.
- LUKEFAHR S., HOHENBOKEN WD., CHEEKE PR., PATTON, NM. 1983a: Doe reproduction and preweaning litter performance of straightbred and crossbred rabbits. *Journal of Animal Science*, **57(5)**: 1090-1099.
- LUKEFAHR S., HOHENBOKEN WD., CHEEKE PR., PATTON NM. 1983b: Characterization of straightbred and crossbred rabbits for milk production and associative traits. *Journal of Animal Science*, **57(5)**: 1100- 1107.
- MASOERO G., UBERTALLE A., MAZZOCCO P., BATTAGLINI LM. 1985: Terminal crossing of New Zealand White and Californian Rabbits. (2) Characteristics on the live animal. *Annalidell Istituto Sperimentale per la Zootecnia*, **18(2)**: 93-109. (*Animal Breeding Abstract*, 55, No. 1264).
- OUDAH SM. 1990: Studies on some rabbit breeds and their crosses. *M. Sci. Thesis, Faculty of Agriculture, Mansoura University, Egypt*.
- OZIMBA CE., LUKEFAHR SD. 1991: Comparison of rabbit breed types of postweaning litter growth, feed efficiency and survival performance traits. *Journal of Animal Science*, **69**: 3494-3500.
- PARTRIDGE GG., FOLEY S., CORRIGALL W. 1981: Reproductive performance in purebred and crossbred commercial rabbits. *Animal Production*, **32**: 325-331.
- ROUVIER R., BRUN JM. 1990: Crossbreeding and selection experiments with rabbits: An overview from studies in France about litter traits. *Options Mediterraneennes, Serie Seminaire*, **8**: 29-34.