



The modulatory impact of Arabic gum and lecithin on the efficiency of cold-stressed Nile tilapia (*Oreochromis niloticus*)

Mohamed R. Soaudy^a, Eman Y. Mohammady^{b,*}, Mohamed A. Elashry^a, Marwa M. Ali^a,
Hoda A.S. Elgarhy^c, Janice Alano Ragaza^d, Mohamed S. Hassaan^{a,*}

^a Department of Animal Production, Fish Research Laboratory, Faculty of Agriculture at Moshtohor, Benha, University, Benha 13736, Egypt

^b Aquaculture Division, National Institute of Oceanography and Fisheries, NIOF, Egypt

^c Department of genetic engineering, Faculty of Agriculture at Moshtohor, Benha, University Benha 13736, Egypt

^d Ateneo Aquatic and Fisheries Resources Laboratory, Department of Biology, School of Science and Engineering, Ateneo de Manila University, Katipunan Ave., Loyola Hts., Metro Manila, Quezon City 1108, Philippines

ARTICLE INFO

Keywords:

Oreochromis niloticus

Lecithin

Winter season

Arabic gum

Cold-stress

ABSTRACT

Tilapia are commonly used in aquaculture but are sensitive to cold, limiting their culture in colder regions. Thus, the purpose of this investigation was to examine how lecithin and/or Arabic gum could help Nile tilapia cope with cold stress during the winter season. In a 3×3 factorial feeding study, tilapia fingerlings with an average initial weight of 7.56 ± 0.10 g were given nine different diets (crude protein: 305 g kg⁻¹; gross energy: 19.47 MJ kg⁻¹) for 60 days during the winter. The diet formulations included three levels of Arabic gum (0 g, 2 g, and 4 g kg⁻¹), with each level receiving three different dosages of lecithin (0 g, 5 g, and 10 g kg⁻¹). After the trial, tilapia fed the food enriched with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin showed the highest specific growth rate, fish survival, weight gain, and the lowest feed conversion ratio. Low-density lipoprotein cholesterol and cholesterol levels were highest in fish given the control diet. The groups fed diets with 10 g kg⁻¹ lecithin and 4 g kg⁻¹ Arabic gum showed higher levels of triglycerides and HDL-C. Significant decreases in alanine aminotransferase, glucose, aspartate aminotransferase, and cortisol activities were also observed ($P < 0.05$) with the same diet. Fish fed a diet enriched with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin showed the highest levels of serum calcium, potassium, sodium, and chlorine, as well as the highest transcription of $\Delta 9D$. The same diet was shown to have the lowest levels of malondialdehyde and the highest activity levels for other antioxidant enzymes such as glutathione peroxidase, superoxide dismutase, glutathione, catalase, and total antioxidant capacity. Tilapia raised in cold-stressed winter conditions had improved survival and performance when fed a diet containing 4 g kg⁻¹ of Arabic gum and 10 g kg⁻¹ of lecithin.

1. Introduction

Physiological stress caused by biological, physical, or chemical stimuli, is the main factor contributing to fish diseases and mortality in aquaculture (Acar et al., 2015; Baba et al., 2016; Hassaan et al., 2019a; Soaudy et al., 2021; Yang et al., 2022; Kesbicç et al., 2022; Mohammady et al., 2023; Elashry et al., 2024; Moawad et al., 2024). Temperature variations are a kind of natural stress that fish may encounter during their life cycle; they are either temporary fluctuations or seasonal changes in fish that are linked to causing disease outbreak and mortality (Ju et al., 2002). Temperature shock can harm fish by disrupting their homeostasis and lowering their metabolic rates (Galloway and Kieffer,

2003; Suski et al., 2006; Vanlandeghem et al., 2010), influencing their swimming ability (Hyvärinen et al., 2004; Suski et al., 2006), weakening immunological syndromes, and increasing opportunistic infections (Chang et al., 2006; Hurst, 2007; Panase et al., 2018). Tropical fish like Nile tilapia optimal temperature range warm water (26–30°C), while cold suboptimal temperatures can decrease feed consumption, growth, and mortality (Bhujel et al., 2007; Azaza et al., 2008; Zerai et al., 2010; Ma et al., 2015; Correa et al., 2017, 2018; Siddik et al., 2014; Nobrega et al., 2017; Wu et al., 2019). Numerous studies have indicated that cold, below-average temperatures have a detrimental effect on Nile tilapia productivity (Ma et al., 2015; Shi et al., 2015; Nobrega et al., 2019; Hassaan et al., 2019a). Therefore, it is critical to increase fish resistance

* Corresponding authors.

E-mail addresses: dreman2529@gmail.com, ey.badiny@niof.sci.eg (E.Y. Mohammady), Mohamed.hassaan@fagr.bu.edu.eg (M.S. Hassaan).

<https://doi.org/10.1016/j.aqrep.2024.102332>

Received 27 February 2024; Received in revised form 27 August 2024; Accepted 31 August 2024

Available online 7 September 2024

2352-5134/© 2024 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC license (<http://creativecommons.org/licenses/by-nc/4.0/>).

to a variety of stressors such as extreme cold (Yokoyama et al., 2005; Shi et al., 2015; Nobrega et al., 2017). Fish, however, can adapt to changes in their immediate environment through altering physiological processes, such as those linked to stress response (Barton, 2002; Siddik et al., 2014).

Various approaches to combat low temperatures include employing greenhouses (Radwan, 2008), using covers made with polyethylene sheets for fish ponds, or utilizing varying pond depths to regulate the water volume (Dan and Little, 2000; Rothbard and Peretz, 2002; Abdel-Aal, 2008). Moreover, techniques for cultivation such as biofloc systems are useful in reducing the negative effects of cold and lowering tilapia mortality over the winter period (Soltan et al., 2015; Soaudy et al., 2021). One promising strategy for preventing environmental stress in aquaculture is the use of functional feed additives (Hassaan et al., 2019b; Bowyer et al., 2019; Refaey et al., 2022; Saleh et al., 2022; Taalab et al., 2022; Abdo et al., 2024), which can help fish that are experiencing growth retardation, inflammatory response, immunosuppression, and oxidative damage. In this context Wu et al., (2019) added *Astragalus membranaceus* extract powder as natural herbs in Nile tilapia diet for recovery cold stress conditions. In cold temperatures, fish increase the unsaturation of fatty acids in their cell membrane phospholipids to maintain fluidity and permeability, aiding in physiological homeostasis (Weber and Bosworth, 2005; Turchini et al., 2009). Numerous fish species have been observed to undergo this process, which is referred to as homeoviscous adaptation (Craig et al., 1995; Fracalossi and Lovell, 1995; Atwood et al., 2003).

Arabic gum is secreted by certain trees, including *Sengalia senegal* and *Acacia senegal*, that contain organic acids, amino acids, a few minerals, and a high-molecular-weight complex of glycoproteins and polysaccharides, such as arabinose, galactose, and rhamnose (Bakhoun et al., 2018; Fouda et al., 2019). Beside the above-mentioned bioactive component in Arabic gum, Imbabi et al. (2023) found another component like, Phosphatidylcholine and phosphatidylserine. The natural prebiotic Arabic gum improves gut health by inhibiting harmful bacteria, promoting nutrient absorption, and activating beneficial bacteria (Al-Baadani et al., 2021). In the study of Naiel et al. (2022), they found that Arabic gum plays a crucial role in promoting antioxidant status, growth, and immunological response of tilapia.

For aquatic species, lecithin is considered a significant source of bioavailable phospholipids (PL). It is an essential component of biological membranes in all eukaryotic cells and plays a crucial function in energy generation through food metabolism (Aničić et al., 2013). In general, diets supplemented with PL resulted in growth-promoting effects and increased survival and stress resistance (Tocher et al., 2008). Adding a lecithin-containing bio-emulsifier to the diet of tilapia has been shown to improve the innate immunological response, growth, and feed digestion of the tilapia (El-Sayed et al., 2021). Furthermore, diets supplemented with marine lecithin (mainly PL) and soy lecithin improved the growth and recovery from stress challenge of juvenile cobia and milkfish larvae (Trushenski et al., 2013; Sivaramakrishnan et al., 2021). Regarding cold stress recovery Batista et al., (2023) found that the addition of soy lecithin in tilapia diet at a range of 42–50 g kg⁻¹ mitigated the cold stress condition. As far as we are aware, there are no scientific reports on the use of Arabic gum or lecithin in tilapia diets as a mitigation feeding strategy during the winter season. Therefore, a feeding trial was done to assess the effects of dietary supplements containing Arabic gum and lecithin on the antioxidant activity, blood index, innate immune responses, and delta-9-desaturase gene (Δ 9D) expression in farmed Nile tilapia during the winter season.

2. Materials and methods

2.1. Determination of bioactive compounds in Arabic gum

Following slight adjustments to the procedure previously described by Folch et al. (1957), the major phospholipids were isolated. Prior to

HPLC analysis, the isolated phospholipid was diluted in a mobile phase solvent that contained 20 % chloroform. With a Porasil silica gel column (10- μ m particle size), an Agilent 1200 Series HPLC system with a computerized solvent supply system and UV detector (Santa Clara, CA, USA) was used to perform the isocratic high-performance liquid chromatographic separation of various phospholipids. The degassed mobile phase, which was supplied at a flow rate of 0.80 mL/min and consisted of acetonitrile, methanol, and 85 % phosphoric acid (96:3:1, v/v/v), was used to elute samples (20 μ L) for HPLC analysis. Using comparable phospholipid standards, the concentration of each sample was determined while the effluent was being observed at a wavelength of 203 nm. The source of the standard (GSH: 200–725–4) was the Sigma Chemical Company located in St. Louis, MO, USA (Fig. 1).

2.2. Design of experiment and diets

Benha University, Faculty of Agriculture's Fish Nutrition Laboratory served as the site of the current experiment. The effects of Arabic gum (Acaci gum; 0 g, 2 g, and 4 g kg⁻¹ diet), lecithin (0 g, 5 g, and 10 g kg⁻¹ diet), and their combinations on the growth, survival rate, serum biochemical parameters, oxidative response, and Δ 9D gene of Nile tilapia fingerlings were investigated in a factorial experiment (3 $\times$ 3). Soybean lecithin was provided by Aceitera Fernal Deheza S. A., Argentina, while Arabic gum was obtained from local market. Nine isonitrogenous (305 g kg⁻¹ crude protein) and isocaloric (19.45 MJ kg⁻¹ gross energy) diets were created by mixing each level of Arabic gum (Table 1). After adding varied amounts of Arabic gum and lecithin, the diet components were thoroughly blended. A pelleting hand-noodle maker was then used to make the 2-mm diameter pellets, which were then left for 12 h at ambient temperature to dry, and finally stored in cellophane bags at -4°C until required, according to Ali et al. (2023). The AOAC (1995) was used to estimate the proximate composition of the diets.

2.3. Fish and experimental management

On 15 November 2021, monosex Nile tilapia, *Oreochromis niloticus* were obtained from the fish farm of the Faculty of Agriculture at Benha University in Egypt. The fish were kept in two cement ponds (2 $\times$ 4 $\times$ 1 m) for 15 days after being collected to acclimate to the experimental conditions and fed a commercial feed (25 g kg⁻¹ protein) before the feeding trial started. Following acclimation, 18 fiberglass tanks (0.5 m³) were randomly filled with 270 uniformly sized fish, with 15 fish per tank as stocking density. The investigation began on 1 December 2021, and it was finished on 1 February 2022 (60 days). The initial body weight of the fish was 7.56 $\pm$ 0.10 g. Two air stones continuously aerated each tank. Fish were manually fed twice daily to maintain the warm water temperature, six days a week (Hassaan et al., 2019a). Table 2 shows the ambient temperature during the experiment period.

Throughout the trial period, water quality data were recorded using various instruments. This included mercury thermometers, an Orion pH meter, and a Jenway 970 dissolved oxygen meter. Standard techniques were used to quantify nitrate (NO₃, 0.55 $\pm$ 0.02 mg L⁻¹), ammonia (NH₄; 0.25 $\pm$ 0.04, mg L⁻¹), and nitrite (NO₂, 0.021 $\pm$ 0.01 mg L⁻¹) once per week according to APHA (1989).

2.4. Growth indices

The growth performance and feed utilization were evaluated by measuring specific growth rate, final body weight, protein efficiency ratio, weight gain, feed conversion ratio, and feed intake as detailed in the footnote of Table 3.

2.5. Serum biochemical analysis

Blood samples were taken from the caudal vein of three fish from




Fig. 1. HPLC chromatogram of Arabic gum. For HPLC analysis, samples (20 mL) were injected and eluted using the degassed mobile phase, which was supplied at a flow rate of 0.80 mL/min and consisted of acetonitrile, methanol, and 85 % phosphoric acid (96:3:1, v/v/v). Using comparable phospholipid standards, the concentration of each sample was determined while the effluent was being monitored at a wavelength of 203 nm. The source of the standard (GSH: 200–725–4) was the Sigma Chemical Company located in St. Louis, MO, USA.

Table 1
Composition and proximate analysis of experimental diets (g kg⁻¹ dry matter).

Ingredient	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6	Diet 7	Diet 8	Diet 9
Fish meal	50	50	50	50	50	50	50	50	50
Soybean meal	450	450	450	450	450	450	450	450	450
Corn gluten	50	50	50	50	50	50	50	50	50
Yellow corn	290	288	286	295	298	291	300	293	296
Wheat bran	100	100	10	100	100	100	100	100	100
Vit&Min ^a	20	20	20	20	20	20	20	20	20
Soyabean oil	40	40	40	30	20	30	20	30	20
Arabic gum	0	2	4	0	2	4	0	2	4
Lecithin	0	0	0	5	10	5	10	5	10
Proximate analysis (%)									
Crude protein	30.02	29.98	29.91	29.88	29.95	30.12	30.10	29.90	30.20
Crude lipid	6.02	6.01	6.02	6.03	6.04	6.01	6.02	6.02	6.04
Ash	4.50	4.40	4.50	4.50	4.50	4.60	4.50	4.50	4.60
Total carbohydrates	59.46	59.61	59.57	59.59	59.51	59.27	59.38	59.58	59.16
Gross energy (MJ kg ⁻¹) ^b	19.45	19.51	19.47	19.49	19.47	19.39	19.40	19.48	19.36

^a Vitamin and mineral mixture kg⁻¹ of mixture contains: 4800 I.U. Vit A, 2400 IU cholecalciferol (vit. D), 40 g Vit E, 8 g Vit K, 4.0 g Vit B12, 4.0 g Vit B2, 6 g Vit B6, 4.0 g, Pantothenic acid, 8.0 g Nicotinic acid, 400 mg Folic acid, 20 mg Biotin, 200 g Choline, 4 g Copper, 0.4 g Iodine, 12 g Iron, 22 g Manganese, 22 g Zinc, 0.04 g Selenium. folic acid, 1.2 mg; niacin, 12 mg; D-calcium pantothenate, 26 mg; pyridoxine. HCl, 6 mg; riboflavin, 7.2 mg; thiamin. HCl, 1.2 mg; sodium chloride (NaCl, 39 % Na, 61 % Cl), 3077 mg; ferrous sulfate (FeSO4.7H2O, 20 % Fe), 65 mg; manganese sulfate (MnSO4, 36 % Mn), 89 mg; zinc sulfate (ZnSO4.7 H2O, 40 % Zn), 150 mg; copper sulfate (CuSO4.5 H2O, 25 % Cu), 28 mg; potassium iodide (KI, 24 % K, 76 % I).

^b Gross energy calculated using gross calorific values of 23.63, 39.52 and 17.15 kJg⁻¹ for protein, fat and carbohydrate, respectively according to Brett (1971).

Table 2
Ambient water temperature average during experimental period from 1 December to 1 February.

Treatment periods	Temperature range	Ambient temperature average (°C)
1 Dec – 14 Dec (1–2 weeks)	16–19.5	18
15 Dec – 29 Dec (2–4 weeks)	14–17	16
30 Dec – 14 Jan (4–6 weeks)	15–18.5	17
15 Jan – 30 Jan (6–8 weeks)	16–21	18.5

each replicate. Enzymes like AST and ALT were measured according to the method of [Reitman and Frankel \(1957\)](#) and serum triglycerides, glucose, LDL-C, and HDL-C were tested by using standardized kits (Modern Laboratory Kits). Cortisol levels were measured using an immulite kit and chemiluminescence, while sodium concentration was determined using a colorimetric detection kit. Potassium, calcium, and

chloride contents were measured using the magon sulfonate method, the cresolphthalein method, and the thiocyanate method, respectively.

2.6. Oxidative enzymes activity

Three fish livers from each replicate were weighed, cleaned, and ground with ice-cold saline (0.1 g of liver in 0.9 mL saline, pH 7.0), then centrifuged at 3000 g for 10 min. The collected supernatant was used to assess the activity of superoxide dismutase (SOD) following [Peskin and Winterbourn \(2000\)](#) approach. For catalase (CAT) activity, [Beers and Sizer's \(1952\)](#) modified procedure was used. Malondialdehyde (MDA) activity was assessed according to [Dogru et al. \(2008\)](#). Glutathione peroxidase (GPx) activity was measured and expressed as units per milligrams of protein following [Moin \(1986\)](#) method. Glutathione (GSH) was measured using the [Beutler et al. \(1963\)](#) technique. The total antioxidant capacity was evaluated using the [Prieto et al. \(1999\)](#) method.

2.7. Expression of delta-9-desaturase gene

Liver samples from fish were taken and homogenized. Total RNA was

Table 3
Growth performance and feed utilization of fish fed supplemented with Arabic gum and lecithin and their interaction.

T	AG ^a g kg ⁻¹ diet	Lecithin g kg ⁻¹	IBW ¹ (g fish ⁻¹)	FBW ² (g fish ⁻¹)	WG ³ (g fish ⁻¹)	SGR ⁴ (% day ⁻¹)	FCR ⁵	FI ⁶ (g fish ⁻¹)	SR ⁷ %
Individual treatments mean									
T1	0	0	7.74	13.11 ^d	5.38 ^e	0.91 ^d	3.46 ^a	18.50 ^b	60.50 ^f
T2	2	0	7.70	18.23 ^c	10.53 ^d	1.49 ^c	2.29 ^b	24.00 ^a	82.50 ^e
T3	4	0	7.67	19.71 ^b	12.04 ^b	1.63 ^b	1.79 ^f	22.50 ^a	85.50 ^d
T4	0	5	7.53	18.97 ^c	11.43 ^c	1.60 ^b	2.06 ^c	23.50 ^a	91.00 ^c
T5	2	5	7.44	18.76 ^c	11.33 ^c	1.60 ^b	1.99 ^d	22.50 ^a	92.50 ^c
T6	4	5	7.44	19.88 ^b	12.45 ^b	1.70 ^b	1.81 ^e	22.50 ^a	98.00 ^{ab}
T7	0	10	7.40	21.78 ^a	14.38 ^a	1.86 ^a	1.60 ^e	23.00 ^a	96.50 ^b
T8	2	10	7.57	20.07 ^{ab}	12.51 ^b	1.68 ^b	1.88 ^e	23.50 ^a	99.00 ^a
T9	4	10	7.54	21.60 ^a	14.07 ^a	1.82 ^a	1.67 ^g	23.50 ^a	99.50 ^a
Pooled SE			0.10	0.58	0.59	0.05	0.23	0.68	1.23
Means of the main effects									
	0		7.61	17.38 ^r	9.77 ^q	1.40 ^q	83.50 ^r	0.29	21.83 ^q
	2		7.51	19.85 ^q	12.33 ^p	1.67 ^p	90.50 ^q	0.27	23.17 ^p
	4		7.55	20.17 ^p	12.63 ^p	1.69 ^p	94.33 ^p	0.35	22.83 ^p
		0	7.70	17.05 ^z	9.35 ^z	1.34 ^y	76.17 ^y	0.31	21.67 ^y
		5	7.47	19.20 ^y	11.73 ^y	1.63 ^x	93.83 ^x	0.35	22.83 ^x
		10	7.50	21.15 ^x	13.65 ^x	1.78 ^x	99.33 ^x	0.26	22.33 ^x
ANOVA (P-value)									
Arabic gum			0.374	0.034	0.027	0.024	0.030	0.2764	0.406
Lecithin			0.353	0.015	0.011	0.008	0.001	0.2166	0.026
Arabic gum × lecithin			0.264	0.001	0.001	0.001	0.001	<.0001	0.011

^aTreatments means represent the average values of three aquaria per treatment. Duncan multiple range test was conducted for individual means only if there was a significant interaction (ANOVA: P < 0.05). Means followed by the same letter are not significantly different.
^bMain effect means followed by the same letter are not significantly different at P < 0.05 by Duncan multiple range test; p, q and r for Arabic gum and x, y and z for lecithin levels.
^c Arabic gum
IBW¹ = initial body weight; FBW² = final body weight; Weight gain (WG)³ = final weight (g) – initial weight (g) Specific growth rate (SGR)⁴: $SGR = \frac{\ln W_2 - \ln W_1}{t} \times 100$ Where: Ln = the natural log W₁ = first fish weight W₂ = the following fish weight in grams t = period in days. Feed conversion ratio (FCR)⁵: $FCR = \frac{\text{Feed ingested (g)}}{\text{Weight gain (g)}}$. FI⁶ = feed intake; SR⁷ = survival fish rate.

isolated from liver samples (three fish from each replicate) by the standard Promega RNA Isolation Kit (Cat No. Z3100, USA) and reagent extraction method according to the manufacturer's instructions. The quantity of the RNA was assessed using a Nano-Drop spectrophotometer (NANODROP 1000, Thermo Scientific, USA). The integrity (quality) was checked by denaturing gel electrophoresis (1 % agarose gel) and the purity by measuring the OD260/OD280 absorption ratio (>1.95).
cDNA was generated from 1 µg of total RNA using High Capacity cDNA (Thermo Fisher Scientific, Cat. No.436, 8814) reverse transcriptase kit for reverse transcriptase polymerase chain reaction (RT-PCR) following the manufacturer's protocol. The product of the first strand cDNA synthesis was stored at -80°C until the quantitative RT-PCR (qRT-PCR) runs.
Delta-9-desaturase (Δ 9D) and 18S rRNA gene primers were purchased from Invitrogen, Germany. The primer sequences of 18S rRNA was F: GTTGCAAAGCTGAAACTTAAAGG; R: TTCCCGTGTGAGTCAAATTAAGC (GenBank no. AF497908.1) and F: ATCACCA-CACGTTCCCATATGAC; R: CCAGACCCAAGAAACATGAAG (GenBank no. AY150696) for Δ 9D. Triplicate qPCR reactions were performed on an AriaMx Real-Time PCR System (Agilent Technologies). Reactions containing 5 µl of 5 × diluted cDNA, 10 pmol each of forward and reverse primers, 0.4 µl ROX dye solution (1:500 dilution) and 10 µl SYBR Green PCR MasterMix (Maxima SYBR Green qPCR, Thermo Fisher Scientific, Cat. No # k0251) were performed in a four-step experimental run protocol: a denaturation program (10 min at 95 °C); an amplification and quantification program repeated 40 times (30 s at 95 °C, 50 s at 55°C and 40 s at 72 °C); a melting curve program (55–95°C with a heating rate of 0.10 °C/s and a continuous fluorescence measurement) and finally a cooling step. Melt curve analyses of the target genes and reference genes resulted in single products with specific melting temperatures. In addition, "no-template" controls (i.e. with water sample) for each set of genes was also run to ensure no contamination of

reagents, no primer-dimer formation. Moreover, 18S rRNA gene was used as an internal standard. The relative mRNA expression levels were calculated by a standard curve method. The expression levels of gene were normalized to the levels of 18S rRNA gene in the same sample. Standard curve was generated by serial dilution of a random mixture of control samples. Relative gene expression ratios (RQ) were calculated using the method of Livak and Schmittgen (2001).
2.8. Data analysis
Before being evaluated, the data were subjected to homogeneity and normality tests. Following that, results were examined using a two-way ANOVA with values for Arabic gum and lecithin. The limit for statistical significance was established at 5 %. Using the SAS ANOVA technique, Duncan's multiple range test was used to determine how the means differed (SAS, version 6.03, Soft Inc., Tusla, OK, USA, SAS, 1993).

3. Results

3.1. Growth and feed indices

The effect of different levels of Arabic gum or lecithin and their interaction in tilapia diet on growth and feed indices are presented in Table 3. Addition of Arabic gum or lecithin individually significantly (P < 0.05) improved final body weight (FBW), survival rate (SR), specific growth rate (SGR), weight gain (WG), and feed conversion ratio (FCR). Positive significant responses were found in WG and SR of tilapia fed diet containing Arabic gum and lecithin (P < 0.05). The fish fed diet with 4 g/kg Arabic gum and 10 g/kg lecithin showed the highest FBW, SGR, WG, and SR, as well as the best FCR.

3.2. Serum biochemical indices

Biochemical indices of fish fed diet supplemented with Arabic gum or lecithin and their interaction are shown in Table 4. Aspartate amino transferase (AST), glucose, alanine aminotransferase (ALT), and cortisol in the blood serum of tilapia were lowered considerably by using Arabic gum or lecithin separately as a dietary supplement ($P < 0.05$). Adding 4 g/kg of Arabic gum and 10 g/kg of lecithin to the fish diet resulted in the lowest levels of ALT, AST, glucose, and cortisol. Individually, there are significant differences in lipid profile including cholesterol, high-density lipoprotein-cholesterol (HDL-C), triglycerides, and low-density lipoprotein-cholesterol (LDL-C) with the addition of Arabic gum and lecithin at varying levels ($P < 0.05$) (Table 5). The lowest values of LDL-C and cholesterol were noted in fish fed control diet, while the highest HDL-C and triglycerides were noted in group fed diet supplemented with 10 g kg⁻¹ lecithin and diet supplemented with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin, respectively

3.3. Blood serum ions

Fish serum ions given a diet enriched with Arabic gum or lecithin and their interaction are presented in Table 6. Sodium, potassium, calcium, and chlorine contents were significantly affected by Arabic gum and lecithin and their interaction. The highest content of sodium, potassium, calcium, and chlorine were obtained by feeding fish a diet enriched with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin.

3.4. Hepatic oxidative biomarkers

Diet supplemented with Arabic gum or lecithin individually significantly ($P < 0.05$) increased the activities of SOD, GPX, CAT, GSH, TAC, and decreased MDA in tilapia (Table 7). The highest CAT, GPX, GSH,

Table 4
Biochemical blood indices of fish fed dietary supplemented with Arabic gum and lecithin and their interaction.

T	AG g kg ⁻¹ diet	Lecithin g kg ⁻¹	Biochemical blood indices			
			ALT (UL ⁻¹)	AST (UL ⁻¹)	Glucose (mg dl ⁻¹)	Cortisol (ng mL ⁻¹)
Individual treatments mean						
T1	0	0	45.25 ^a	26.90 ^a	119 ^a	14.5 ^a
T2	2	0	40.25 ^b	24.65 ^b	111 ^b	12.5 ^{cb}
T3	4	0	38.75 ^c	23.10 ^c	101.5 ^c	12.65 ^{cb}
T4	0	5	38.10 ^d	21.25 ^d	91.5 ^d	12.5 ^{cb}
T5	0	10	34.10 ^g	13.90 ^g	73.5 ^g	13 ^b
T6	2	5	37.40 ^e	19.65 ^e	87 ^e	11.5 ^d
T7	2	10	33.10 ^h	13.15 ^h	69 ^h	12.25 ^c
T8	4	5	35.25 ^f	14.65 ^f	81.5 ^f	12.25 ^c
T9	4	10	31.10 ⁱ	12.25 ⁱ	62.5 ⁱ	10.75 ^e
Pooled SE			0.17	0.17	0.36	0.14
Means of the main effects						
	0		391.5 ^p	206.83 ^p	94.67 ^p	13.33 ^p
	2		369.17 ^q	191.50 ^q	89.00 ^q	12.08 ^q
	4		350.33 ^r	166.67 ^r	81.83 ^r	11.88 ^q
		0	414.17 ^x	248.83 ^x	110.50 ^x	13.22 ^x
		5	369.17 ^y	185.17 ^y	86.67 ^y	12.08 ^y
		10	327.67 ^z	131 ^z	68.33 ^z	12 ^y
ANOVA (P-value)						
Arabic gum			0.0001	0.0002	0.1133	0.0262
Lecithin			0.0001	0.0001	0.0725	0.0031
Arabic gum × Lecithin			0.0001	0.0001	0.0001	0.0001

[†]Treatments means represent the average values of three aquaria per treatment. Duncan multiple range test was conducted for individual means only if there was a significant interaction (ANOVA: $P < 0.05$). Means followed by the same letter are not significantly different.

[‡]Main effect means followed by the same letter are not significantly different at $P < 0.05$ by Duncan multiple range test; p, q and r for Arabic gum and x, y and z for lecithin levels.

[‡] Arabic gum

Table 5

Serum lipid profile of fish fed dietary supplemented with Arabic gum and lecithin and their interaction.

T	AG g kg ⁻¹ diet	Lecithin g kg ⁻¹	Lipid profile			
			Cholesterol (mmol L ⁻¹)	Triglyceride (mmol L ⁻¹)	HDL (mmol L ⁻¹)	LDL (mmol L ⁻¹)
Individual treatments mean						
T1	0	0	119.00 ^h	202.50 ^f	44.50 ^d	49.00 ^e
T2	2	0	186.00 ^f	216.50 ^e	29.50 ^g	69.01 ^a
T3	4	0	175.50 ^g	180.50 ^g	34.00 ^f	59.01 ^c
T4	0	5	211.50 ^e	178.50 ^g	44.50 ^d	63.50 ^b
T5	0	10	401.00 ^a	331.50 ^a	52.50 ^b	52.00 ^d
T6	2	5	316.50 ^c	281.00 ^c	36.00 ^f	42.50 ^f
T7	2	10	399.00 ^a	299.00 ^b	49.02 ^c	68.51 ^a
T8	4	5	311.00 ^d	201.00 ^f	41.00 ^e	52.02 ^d
T9	4	10	387.00 ^b	273.00 ^d	59.00 ^a	64.00 ^b
Pooled SE			1.31	0.86	0.87	0.42
Means of the main effects						
	0		243.83 ^q	237.50 ^{pq}	47.00 ^p	54.83
	2		300.50 ^p	265.50 ^p	38.17 ^q	60.00
	4		291.17 ^p	218.17 ^q	44.67 ^p	58.33
		0	160.17 ^z	199.83 ^y	35.83 ^z	59.00
		5	279.67 ^y	220.17 ^y	40.50 ^y	52.67
		10	395.67 ^x	301.17 ^x	53.50 ^x	61.50
ANOVA (P-value)						
Arabic gum			0.0063	0.0400	0.0032	0.6327
Lecithin			0.0001	0.0001	0.0001	0.2800
Arabic gum× Lecithin			0.0001	0.0001	0.0001	0.0001

[†]Treatments means represent the average values of three aquaria per treatment. Duncan multiple range test was conducted for individual means only if there was a significant interaction (ANOVA: $P < 0.05$). Means followed by the same letter are not significantly different.

[‡]Main effect means followed by the same letter are not significantly different at $P < 0.05$ by Duncan multiple range test; p, q and r for Arabic gum and x, y and z for lecithin levels.

[‡] Arabic gum

SOD, and TAC were recorded in fish fed dietary supplemented with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin, whereas fish fed the same supplements had the lowest levels of MDA.

3.5. Gene expression

Fig. 2 shows the effect of the different levels of Arabic gum or lecithin on the expression of the delta-9-desaturase (Δ 9D) gene in Nile tilapia livers. The transcription of Δ 9D was upregulated in groups fed with diets supplemented with the test ingredients than those fed diets without supplementation. Also, fish fed Arabic gum levels showed significantly higher Δ 9D expression compared with those fed lecithin levels. The highest transcription of Δ 9D was seen in fish given diet that included 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin.

4. Discussion

4.1. Growth performance and survival

Cold stress is one of the main hazards in the aquaculture production industry. It results in physiological dysfunction and has a detrimental impact on fish performance and survival (Bhujel et al., 2007; Nobrega et al., 2017; Cheng et al., 2018; Hassaan et al., 2019a). The current study shown that adding 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin to the feed could reduce the effects of cold stress and increase the survival of Nile tilapia. Very little research has looked at the potential effect of dietary Arabic gum usage on fish performance (Faggio et al., 2015; El-faki, 2016; Naiel et al., 2022). Moreover, there is no information available on the use of Arabic gum to reduce cold stress in Nile tilapia. The improvement in survival of Nile tilapia in the present study may be due to bioactive compounds in Arabic gum (Fig. 1). Naiel et al. (2022)

Table 6
Serum blood ions of fish fed dietary supplemented with Arabic gum and lecithin and their interaction.

T	AG g kg ⁻¹ diet	Lecithin g kg ⁻¹	Blood ions			
			Na (mEq l ⁻¹)	K (mEq l ⁻¹)	Ca (mg dl ⁻¹)	Cl (mEq l ⁻¹)
Individual treatments mean						
T1	0	0	139.00 ^e	2.30 ^d	9.13 ^d	91.50 ^f
T2	2	0	142.00 ^d	2.90 ^c	11.10 ^c	111.00 ^e
T3	4	0	146.15 ^c	3.20 ^b	12.30 ^b	113.00 ^d
T4	0	5	148.50 ^c	3.10 ^b	11.60 ^c	112.00 ^d
T5	0	10	153.00 ^b	3.50 ^b	12.90 ^b	120.00 ^b
T6	2	5	152.00 ^b	3.90 ^b	12.70 ^b	122.00 ^a
T7	2	10	149.00 ^c	3.20 ^b	11.05 ^c	110.00 ^e
T8	4	5	147.40 ^c	3.80 ^b	12.90 ^b	118.00 ^c
T9	4	10	166.00 ^a	4.20 ^a	13.20 ^a	123.00 ^a
Pooled SE			0.34	0.83	0.19	0.71
Means of the main effects						
	0		145.50 ^f	2.87 ^f	10.59 ^f	104.50 ^f
	2		147.47 ^q	3.40 ^q	12.30 ^q	116.33 ^q
	4		154.72 ^p	3.77 ^p	12.73 ^p	119.33 ^p
		0	142.38 ^z	2.80 ^z	10.84 ^y	105.17 ^y
		5	151.17 ^y	3.50 ^y	12.40 ^x	118.00 ^x
		10	154.72 ^x	3.77 ^x	12.73 ^x	119.33 ^x
ANOVA (P-value)						
Arabic gum			0.4277	0.0304	0.0	0.0217
Lecithin			0.0576	0.0388	0.0304	0.0159
Arabic gum× Lecithin			0.0001	0.0195	0.0004	0.0001

[†]Treatments means represent the average values of three aquaria per treatment. Duncan multiple range test was conducted for individual means only if there was a significant interaction (ANOVA: P < 0.05). Means followed by the same letter are not significantly different.

[‡]Main effect means followed by the same letter are not significantly different at P < 0.05 by Duncan multiple range test; p, q and r for Arabic gum and x, y and z for lecithin levels.

[§] Arabic gum

detected natural chemicals such as glucuronic acid, polysaccharides, minerals, and neutral sugars (e.g., rhamnose, arabinose, and galactose) in Arabic gum. These compounds are reported to impart

anti-inflammatory, antioxidant, and immunostimulatory effects in fish. Furthermore, earlier research showed that Arabic gum is a natural prebiotic that stimulates growth (Calame et al., 2008). Arabic gum contains polysaccharides, glucuronic acid, and minerals that promote performance by fostering the growth of beneficial bacteria (Al-Baadani et al., 2021).

On the other hand, lecithin plays an important role in energy production through nutritional metabolism, notably fat digestion and absorption via emulsification and facilitation of the storage of lipids in the hepatopancreas; it is also a necessary part of the biological membranes of all eukaryotic cells (Hamre et al., 2013). The feed digestion, growth, and innate immune status of tilapia were all significantly strengthened by the dietary addition of bioemulsifiers including lecithin (El-Sayed et al., 2021). Lecithin also plays a cooperative function in the intestinal absorption of cholesterol, which enhances animal development and survival (Coutteau et al., 1997; Haghparast et al., 2019). Furthermore, phospholipids have growth-promoting, antioxidant, and chemo-attractant qualities that enhance resilience to stress, particularly in the early stages of development (Hertrampf and Piedad-Pascual, 2000; Tocher et al., 2008; El-Naggar et al., 2021). Dietary lecithin increases lipoprotein synthesis and secretion as well as enhances feed digestibility. Thus, fish somatic development is improved using dietary lipids (Seiliez et al., 2006). The findings of the current study on improved development and survival may be due to lecithin though its crucial function in energy production through nutritional metabolism, particularly fat digestion and absorption via emulsification (Gisbert et al., 2005; Morais et al., 2007).

4.2. Serum biochemical parameters and lipid metabolism

Biochemical blood indices can indicate the health of fish. Monitoring the biochemical fluctuations in hepatic enzyme activities, such as ALT and AST can provide insight into liver function (Zhai et al., 2014; Hassaan et al., 2015; Hassaan et al., 2019b; Hassaan et al., 2021; Inanan et al., 2021; Alandiyyany et al., 2022; Navruz et al., 2023; Awad et al., 2024). It well known that the activities of aminotransferase enzymes are

Table 7
Hepatic oxidative response (U/g protein) of fish fed dietary supplemented with Arabic gum and lecithin and their interaction.

T	AG g kg ⁻¹ diet	Lecithin g kg ⁻¹	Antioxidant enzymes					
			SOD	CAT	GPX	GSH	MDA	TAC
Individual treatments mean								
T1	0	0	586.21 ^f	478.60 ^d	99.00 ^g	49.00 ^g	50.33 ^a	4.62 ^d
T2	2	0	759.29 ^e	488.60 ^d	121.00 ^f	70.03 ^f	33.47 ^b	7.05 ^c
T3	4	0	1005.20 ^b	510.00 ^e	131.00 ^e	73.00 ^e	29.00 ^b	8.00 ^c
T4	0	5	903.44 ^c	512.00 ^c	146.00 ^d	75.00 ^e	19.27 ^c	8.20 ^c
T5	0	10	992.23 ^c	540.00 ^b	181.50 ^b	89.50 ^d	12.00 ^d	9.30 ^b
T6	2	5	892.47 ^d	562.00 ^a	161.50 ^d	91.00 ^d	14.00 ^d	10.30 ^b
T7	2	10	946.34 ^c	511.00 ^c	190.50 ^a	110.00 ^c	7.55 ^e	9.30 ^b
T8	4	5	1106.45 ^b	567.00 ^a	171.50 ^c	119.00 ^b	7.00 ^e	11.80 ^a
T9	4	10	1129.40 ^a	570.00 ^a	188.00 ^a	128.00 ^a	6.10 ^f	12.90 ^a
Pooled SE			1.09	8.91	0.79	0.84	0.89	0.55
Means of the main effects								
	0		812.00 ^f	500.53 ^f	145.17 ^f	78.00 ^f	25.72 ^f	7.37 ^f
	2		952.66 ^q	531.87 ^q	158.00 ^q	92.84 ^q	17.49 ^q	9.38 ^q
	4		1009.02 ^p	547.33 ^p	160.17 ^p	97.33 ^p	16.37 ^p	10.40 ^p
		0	783.57 ^z	492.40 ^z	117.00 ^z	64.01 ^z	37.60 ^x	6.56 ^z
		5	929.38 ^y	538.00 ^y	163.00 ^y	85.17 ^y	15.09 ^y	9.27 ^y
		10	1060.73 ^x	549.33 ^x	183.33 ^x	119.00 ^x	6.88 ^z	11.33 ^x
ANOVA (P-value)								
Arabic gum			0.0001	0.0424	0.0001	0.0001	0.0109	0.04877
Lecithin			0.0001	0.0285	0.0001	0.0001	0.0001	0.0487
Arabic gum× Lecithin			0.0001	0.0001	0.0001	0.0001	0.0001	0.0144

[†]Treatments means represent the average values of three aquaria per treatment. Duncan multiple range test was conducted for individual means only if there was a significant interaction (ANOVA: P < 0.05). Means followed by the same letter are not significantly different.

[‡]Main effect means followed by the same letter are not significantly different at P < 0.05 by Duncan multiple range test; p, q and r for Arabic gum and x, y and z for lecithin levels.

[§] Arabic gum

Two way ANOVA (P-Value): Arabic gum (AG) = 0.023; Lecithin (L) = 0.0035; Arabic gum $\times$ Lecithin (AG $\times$ L) = 0.002




Fig. 2. Gene expression of delta-9-desaturase in liver of Nile tilapia fingerlings fed diet supplemented with different levels of Arabic gum and lecithin during the winter season. T1 = (0 g AG + 0 g kg⁻¹ diet); T2 = (2 g AG + 0 g kg⁻¹ diet); T3 (4 g AG + 0 g kg⁻¹ diet); T4 = (0 g AG + 5 g kg⁻¹ diet); T5 (0 g AG + 10 g kg⁻¹ diet); T6 = (2 g AG + 5 g kg⁻¹ diet); T7 = (2 g AG + 10 g kg⁻¹ diet); T8 = (4 g AG + 5 g kg⁻¹ diet); T9 = (4 g AG + 10 g kg⁻¹ diet).

sensitive to cold temperature. Previous studies found that higher activities of ALT and AST when fish are exposed to low temperatures (Shi et al., 2015; Panase et al., 2018; Wu et al., 2019; Hassaan et al., 2019a; Soaudy et al., 2021). As far as we know, there have been no studies on the effect of dietary Arabic gum and lecithin on the activities of aminotransferase enzymes and serum biomarkers of stress in fish under cold stress conditions. While, there are other feed additives improved the levels of ALT and AST such as chia seed powder when added in tilapia diet significantly decreased the levels of ALT and AST (Mahmoud et al., 2023). In the present study dietary Arabic gum and lecithin at level 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin improved AST and ALT activities of Nile tilapia reared in cold stress. This finding may indicate the hepatoprotective role of Arabic gum and lecithin. While, rise of ALT and AST might be a sign of increased hepatic enzyme production or of enzymes leaking via harmed plasma membranes (Yang and Chen, 2003). The reduction of ALT and AST in tilapia fed diet supplemented with Arabic gum and lecithin in the present study reflect the positive effect of the diet additives for cold stress recovery. In line with the present study Naiel et al., (2022) found that addition of Arabic gum at 1 % in tilapia diet resulted in decreased ALT and AST activities. Furthermore, lecithin could improve the metabolism activities and maintain health status of fish (Attia and Kamel, 2012; Shang et al., 2018).

Complex hormonal interactions, including those involving cortisol, regulate the amount of glucose in the blood (Pacheco and Santos, 2001; Panase et al., 2018). Cortisol regulates glucose mobilization through gluconeogenesis and is considered an indicator of stress (Vijayan et al., 1997; Wendelaar Bonga, 1997; Mommsen et al., 1999). Fish use glucose as a fuel source that is stored as glycogen in their muscles and liver. Compared to serum cholesterol and protein, it is more susceptible to temperature variations (Lucas, 1996; Lermen et al., 2004). The current study found that dietary supplemented with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin decreased the levels of cortisol and glucose of tilapia serum compared to the other treatments. While, increased cortisol and glucose levels in fish fed diet devoid of Arabic gum and/or lecithin may be the cause of the increased energy need to combat cold stress and compensate for the decreased enzyme response (Atwood et al., 2003). Correspondingly with the present data, Tandon and Joshi (1974), Best et al. (2001), and Cho et al. (2015) reported that cold stress caused elevated blood glucose levels in *Clarias batrachus*, *Salmo trutta*, red spotted grouper, and *Epinephelus akaara*, resulting in retarded fish metabolism. Our data indicates that lecithin or Arabic gum supplements have an anti-stress effect. Fish given a diet containing 4 g kg⁻¹ Arabic

gum and 10 g kg⁻¹ lecithin had lower levels of cortisol and glucose. According to Segvić-Bubić et al. (2013), adding 2.5 g kg⁻¹ of propolis to diets could help prevent oxidative stress in sea bass when exposed to short-term (24-hour) cold stress. This is based on the impact of propolis on the levels of serum cortisol and glucose in fish experiencing short-term cold stress. Additionally, according to Acar (2018), Nile tilapia reared under standard rearing conditions and given varied doses of propolis extract showed no appreciable variations (P > 0.05) in glucose content. Further studies needed to clarify the role of bioactive compounds to decrease the serum biochemical parameters as a biomarker.

LDL-C carries cholesterol from the liver to tissues, while HDL-C carries cholesterol from peripheral tissues into hepatic cells for additional metabolism (Jafari et al., 2018). Elevated triglyceride (TG) levels can result from glycogen storage or nephrotic syndrome. Monitoring these levels helps assess lipid metabolism, nutritional status, and liver health (Coz-Rakovac et al., 2005; Osman et al., 2010). The current study showed high triglyceride and cholesterol levels in fish fed diets supplemented with 10 g kg⁻¹ lecithin than diets free of the test ingredients. This could suggest the capacity of lecithin to lessen the cold stress response of tilapia. The breakdown of TG to create adenosine triphosphate may cause a decrease in TG levels in the diet without lecithin (Sun et al., 2019). This can interfere with the liver's ability to mobilize fat, alter the fluidity of membranes, and affect how the liver and tissues circulate blood in response to cold stress (Chang et al., 2006). Also, Jafari et al. (2018) found that higher cholesterol in fish fed dietary supplemented with soya lecithin, due to higher hepatic synthesis of cholesterol. Also, the present study showed that the level of HDL-C content tends to increase in the serum of fish fed dietary supplemented with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ diet lecithin, while LDL-C increased in fish fed 2 g kg⁻¹ Arabic gum followed by fish fed 2 g kg⁻¹ diet Arabic gum and 10 g kg⁻¹ lecithin. The rise in HDL-C indicates that it is transferring cholesterol from peripheral cells to the liver to reduce damage from lipid peroxidation due to cold stress (Fredenrich and Bayer, 2003). Higher levels of blood triglycerides, LDL-C, cholesterol, and HDL-C herein suggest that lecithin and Arabic gum are beneficial in avoiding liver tissue damage under cold stress circumstances. In a study by Segvić-Bubić et al. (2013), there was a decrease in triglyceride levels in *Dicentrarchus labrax* before and after experiencing acute low-temperature stress for 24 h. Furthermore, propolis extract in varying amounts was added to the Nile tilapia diets and showed no significant variations in their triglyceride levels (Acar, 2018). According

to Hassaan et al. (2019a), tilapia fed diets with propolis extract tend to increase blood HDL-C, triglyceride, LDL-C, and cholesterol contents at a faster pace than fish fed diets without the supplement. Additionally, a study on *Sparus aurata*, revealed that serum triglyceride contents rose following cold stress, indicating enhanced energy requirements to deal with the restoration processes as shown with the use of triglycerides (Kyprianou et al., 2010). The disagreement between our results and others could be caused by variations in the species, exposure period to cold stress, and water temperature.

4.3. Blood serum ions

Cold stress conditions adversely alter the plasma electrolytes stress (Cerqueira and Fernandes, 2000). Trophic fish blood osmolality stress indicators include plasma levels of Ca. Weyh et al. (2022) reported that minerals play a crucial role in regulating inflammation and supporting immune function. The present study showed that the blood ion concentrations of Na, Ca, Cl, and K were higher in fish that received nutritional supplements containing 4 g kg⁻¹ of Arabic gum and 10 g kg⁻¹ of lecithin in diet. This may be because Arabic gum and lecithin have a beneficial effect on regulating blood osmotic pressure during cold temperature stress (Tyagi et al., 2013; Soaudy et al., 2021). The concentration of K and Na ions in the plasma was shown to be crucial for the survival of *Shizothorax richardsonii* at low temperatures, as supported by our observations. The levels of Na and K in fish fed control diet decreased during cold stress, according to Hassaan et al. (2019a), but their levels in fish fed diets supplemented with propolis extract increased significantly. Stress and disturbances can alter electrolyte levels (such as K and Na) and osmoregulation by activating the pituitary-internal axis in the neuroendocrine system, leading to the release of stress hormones like corticosteroids and catecholamines.

4.4. Hepatic oxidative stress

Exposure to cold temperature stress alters antioxidant capacity, which leads to an increase in energy metabolism (Ye et al., 2015; Kaushik and Kaur, 2003). The liver contains antioxidant enzymes that are capable of scavenging free radicals such as hydroxyl radicals and superoxide anions to avoid being damaged by oxidation and shield the body from harm (Lortz et al., 2000; Zikic et al., 2001; Rudneva et al., 2010; Sandamalika et al., 2021; Mohammady et al., 2024). Lipid peroxidation produces MDA, which can be elevated to the point that it inactivates enzymes and decreases the permeability of cell membranes, causing oxidative damage to proteins, DNA, and cytoplasm, particularly when fish are exposed to unfavorable climatic circumstances (Tüzgen et al., 1998; Yao et al., 2010; Garcia et al., 2020). The current investigation revealed increased CAT, GSH, SOD, and TAC activities in tilapia fed a diet enriched with 4 g kg⁻¹ of Arabic gum with 10 g kg⁻¹ lecithin. The addition of Arabic gum and lecithin to the fish diets may result in improved antioxidant defense when exposed to cold stress. In this context, Naiel et al. (2020) found that diets supplemented with Arabic gum improved the antioxidant enzymes activities of GSH, SOD, and CAT, and decreased MDA of tilapia. Moreover, Arabic gum can enhance the antioxidant activity of liver cells by regulating the expression of genes related to oxidative stress (Musa et al., 2013). Also, Mirghani et al. (2018) found that the potent antioxidant properties of Arabic gum may be attributed to its high concentration of phenolic compounds. Furthermore, Arabic gum has high antioxidant activity and a protective effect against oxidative damage and a slight immunostimulatory effect when added as food supplement in fish diet. In the study by Kumar et al. (2014), lecithin was found to have a positive effect on the liver's GSH, CAT, and SOD in milkfish at different temperatures. It also protected cells from stress-induced damage, as supported by the current data.

4.5. Gene expression

Certain genes, such as delta 9 desaturase (Δ 9D), are highly expressed in a variety of poikilothermic species throughout the cold acclimation process and vary in expression to adapt to the cold (Zeraf et al., 2010; Hassaan et al., 2019; Wu et al., 2020). According to earlier research, temperature, feed quality, and the various fatty acid types have an impact on the expression of the Δ 9D gene (Tocher et al., 1996; Hsieh et al., 2007). Desaturation membrane lipids is one of Δ 9D primary roles in maintaining membrane fluidity in the cold. Our results showed that fish fed diets with Arabic gum and lecithin had enhanced mRNA transcriptome of the Δ 9D gene in liver tissue compared to fish fed control diet under cold stress (Fig. 2). The fish liver tissue samples supplemented with 4 g kg⁻¹ of Arabic gum and 10 g kg⁻¹ of lecithin showed the highest Δ 9D expression. Low temperatures enhance the fluidity of the cell membrane by increasing the unsaturation of phospholipids more than high temperatures (Hagar and Hazel, 1985b; Ruyter et al., 2003; Soaudy et al., 2021). Decreased expression of genes for acetyl-CoA carboxylase, fatty acid synthase, and carnitine palmitoyl transferase indicates the harmful effects of cold stress, such as inhibition of lipid synthesis, digestion, and oxidation. Conversely, there was a rise in the need of *Cherax quadricarinatus* for highly unsaturated fatty acids due to elevated expression of the sphingolipid delta-4 desaturase gene under cold stress (Wu et al., 2020). The decline in fish survival herein when fed diets without supplementation might have prevented the phosphatidylcholine from being replaced by phosphatidylinositol in cell membranes, which might have also made it more difficult for the fish to withstand the cold (Greene and Selivonchick, 1987; Henderson and Tocher, 1987). This may not be solely due to cold shock, but rather a decrease in metabolic rate caused by reduced feed consumption affecting liver metabolism. Furthermore, under optimal temperature conditions, Δ 9D expression was considerably higher in hybrid tilapia (*Oreochromis niloticus* $\times$ *O. aureus*) fed diet containing coconut oil than in those provided diets containing palmitoleic or fish oil (Hsieh et al., 2007). The amount of unsaturated fat in rainbow trout liver can be changed by modifying diets, according to Tocher et al. (1996). The current study found that Arabic gum levels had significantly higher Δ 9D expression compared to lecithin levels due to higher levels of phosphatidylcholine and phosphatidylserine. This finding is intriguing and requires further investigation.

5. Conclusion

The feeding of Nile tilapia supplemented with 4 g kg⁻¹ Arabic gum and 10 g kg⁻¹ lecithin has the potential to improve fish physiological status, reduce mortality, and boost resilience against harsh wintertime temperatures. These results could be applied in the aquatic feed industry as additives for winter diets. New feed technologies are necessary to increase the tilapia output in subtropical regions. To increase production in these areas, research on winter feed designs, sustainable husbandry practices, and cold-resistant Nile tilapia breeds are recommended.

Ethical approval

All experiments were approved by the authority of NIOF Committee for Institutional Care of Aquatic Organisms and Experimental Animals (NIOF-AQ4-F-23-R-051).

CRediT authorship contribution statement

Mohamed Elashry: Data curation. **Jiance Ragaza:** Writing – review & editing. **Eman mohammady:** Writing – review & editing, Writing – original draft, Software, Methodology. **Mohamed R. Soaudy:** Conceptualization. **Mohamed Hassaan:** Writing – review & editing, Writing – original draft, Software. **Janice Alano Ragaza:** Funding acquisition. **Hoda Elgarhy:** Formal analysis. **Marwa M. Ali:** Data curation.

Declaration of Competing Interest

None

Data Availability

Data will be made available on request.

Acknowledgments

The authors would like to thank the National Institute of Oceanography and Fisheries (NIOF), Egypt and Benha University for their cooperation. The authors are also thankful to the Ateneo de Manila University for the publication aid and support.

References

- Abdel-Aal, M.M., 2008. Effects of over-wintering and water depth on growth performance of Nile tilapia (*Oreochromis niloticus*). Proceedings of the 8th International Symposium on Tilapia in Aquaculture. The Central Laboratory for Aquaculture Research, Ministry of Agriculture and Land Reclamation, Cairo, pp. 297–306.
- Abdo, H.S., Mohammady, E.Y., Tonsy, Hayam D., Ibrahim, H.D., Hassaan, A.-K., M.S., 2024. The potential synergistic action of quercetin and/or *Pediococcus acidilactici* on Nile tilapia, *Oreochromis niloticus* performance. Aquaculture 581, 740353.
- Acar, U., 2018. Effects of diet supplemented with ethanolic extract of propolis on growth performance, hematological and serum biochemical parameters and disease resistance of Mozambique tilapia (*Oreochromis mossambicus*) against *Streptococcus iniae*. Aquaculture 495, 339–344.
- Acar, Ü., Kesbiç, O.S., Yilmaz, S., Gültepe, N., Ali Türker, A., 2015. Evaluation of the effects of essential oil extracted from sweet orange peel (*Citrus sinensis*) on growth rate of tilapia (*Oreochromis mossambicus*) and possible disease resistance against *Streptococcus iniae*. Aquaculture 437, 282–286.
- Alandiyjany, M.N., Kishawy, A.T.Y., Abdelfattah-Hassan, A., Eldoumani, H., Elazab, S.T., El-Mandrawy, S.A.M., Ibrahim, D., 2022. Nano-silica and magnetized-silica mitigated lead toxicity: their efficacy on bioaccumulation risk, performance, and apoptotic targeted genes in Nile tilapia (*Oreochromis niloticus*). Aquat. Toxicol. 242, 106054.
- Al-Baadani, H.H., Al-Mufarrej, S.I., Al-Garadi, M.A., Alhidary, I.A., Al-Sagan, A.A., Azzam, M.M., 2021. The use of gum Arabic as a natural prebiotic in animals: a review. Anim. Feed Sci. Technol., 114894.
- Ali, M.M., Elashry, M.A., Mohammady, E.Y., Soaudy, M.R., El-Garhy, H.S., El-Erian, M. A., Mustafa, Ahmed, Abouelsoud, Mohamed, Ragaza, Janice A., El-Haroun, Ehab R., Hassaan, M.S., 2023. Dietary Alpha-Monolaurin for Nile Tilapia (*Oreochromis niloticus*): stimulatory effects on growth, immunohematological indices, and immune-related gene expressions. Aquac. Res. 2023, 3155447 <https://doi.org/10.1155/2023/3155447>.
- Aničić, I., Treer, T., Matulić, D., Safner, R., Tomljanović, T., Piria, M., Šprem, N., 2013. Effects of dietary vitamin C and soybean lecithin in the nutrition of brown bullhead (*Ameiurus nebulosus* L.) fingerlings. Ital. J. Anim. Sci. 12 (2), e27 <https://doi.org/10.4081/ijas.2013.e27>.
- AOAC, 1995. Association Of Official Analytical Chemists. In: Cunni, P.A. (Ed.), Official Methods of Analysis of the Association Official Analytical Chemists, 16th. 1. AOAC International, Arlington, USA, pp. 1298.
- APHA, (American Public Health Association), 1989. Water Pollution Control Federation. Standard methods for the examination of water and wastewater 17.
- Attia, Y.A., Kamel, K.I., 2012. Semen quality, testosterone, seminal plasma biochemical and antioxidant profiles of rabbit bucks fed diets supplemented with different concentrations of soybean lecithin. Animal 6 (5), 824–833. <https://doi.org/10.1017/S1751731111002229>.
- Atwood, H.L., Tomasso, J.R., Webb, K., Gatlin III, D.M., 2003. Low-temperature tolerance of Nile tilapia, *Oreochromis niloticus*: effects of environmental and dietary factors. Aquac. Res. 34 (3), 241–251.
- Awad, A.-H., Mohammady, E.Y., Soaudy, M.R., Rabetimarghezar, N., El-Haroun, E.R., Hassaan, M.S., 2024. Growth and physiological response of Nile tilapia (*Oreochromis niloticus*) fed a fermented mixture of plant protein sources. Anim. Feed Sci. Technol. 315, 116034.
- Azaza, M.S., Dhraief, M.N., Kraiem, M.M., 2008. Effects of water temperature on growth and sex ratio of juvenile Nile tilapia *Oreochromis niloticus* (Linnaeus) reared in geothermal waters in southern Tunisia. J. Therm. Biol. 33 (2), 98–105.
- Baba, E., Acar, Ü., Ontas, C., Kesbiç, O.S., Yilmaz, S., 2016. The use of Avena sativa extract against *Aeromonas hydrophila* and its effect on growth performance, hematological and immunological parameters in common carp (*Cyprinus carpio*). Ital. J. Anim. Sci. 15 (2), 325–333.
- Bakhoun, N., Fall, D., Fall, F., Diouf, F., Hirsch, A., Balachandrar, D., Diouf, D., 2018. *Senegalia senegal* (synonym: *Acacia senegal*), its importance to sub-Saharan Africa, and its relationship with a wide range of symbiotic soil microorganisms. South Afr. J. Bot. 119, 362–368.
- Barton, B.A., 2002. Stress in fishes: a diversity of responses with particular reference to changes in circulating corticosteroids. Integr. Comp. Biol. 42 (3), 517–525.
- Batista, R.O., Richter, B.L., Banze, J.F., Schleider, D.D., Salhi, M., Nobrega, R.O., Silva, M. F.O.D., Mattioni, B., Pettigrew, J.E., Fracalossi, D.M., 2023. Soy Lecithin supplementation promotes growth and increases lipid digestibility in GIFT Nile Tilapia raised at suboptimal temperature. Fishes 8 (8), 404.
- Beers, R.F., Sizer, I.W., 1952. A spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. J. Biol. Chem. 195, 133–140.
- Best, J.H., Eddy, F.B., Codd, G.H., 2001. Effects of purified microcystin-LR and cell extracts of Microcystis strains PCC 7813 and function in brown trout (*Salmo trutta*) alevins. Fish. Physiol. Biochem. 24, 171–178.
- Beutler, E., Duron, O., Kelly, B.M., 1963. Improved method for the determination of blood glutathione. J. Lab. Clin. Med. 61, 882–890.
- Bhujel, R.C., Little, D.C., Hossain, A., 2007. Reproductive performance and the growth of pre-stunted and normal Nile tilapia (*Oreochromis niloticus*) broodfish at varying feeding rates. Aquaculture 273 (1), 71–79.
- Bowyer, P.H., El-Haroun, E.R., Hassaan, M., Salim, H., Davies, S.J., 2019. Dietary nucleotides enhance growth performance, feed efficiency and intestinal functional topography in European Seabass (*Dicentrarchus labrax*). Aquac. Res. 50 (7), 1921–1930.
- Calame, W., Weseler, A.R., Viebke, C., Flynn, C., Siemensma, A.D., 2008. Gum Arabic establishes prebiotic functionality in healthy human volunteers in a dose-dependent manner. Br. J. Nutr. 100, 1269–1275.
- Cerqueira, C.C.C., Fernandes, M.N., 2000. Gill tissue recovery after copper exposure and blood parameter responses in the tropical fish *Prochilodus scrofa*. Ecotoxicol. Environ. Saf. 52, 83–91.
- Chang, Y.M., Cao, D.C., Sun, X.W., Liang, L.Q., 2006. Changes of serum biochemical indices of common carp affected by cold temperatures. Chin. J. Fish. 2, 71–75.
- Cheng, C.-H., Guo, Z.-X., Wang, A.-L., 2018. The protective effects of taurine on oxidative stress, cytoplasmic free-Ca²⁺ and apoptosis of pufferfish (*Takifugu obscurus*) under low temperature stress. Fish. Shellfish Immunol. 77, 457–464. <https://doi.org/10.1016/j.fsi.2018.04.022>.
- Cho, H.C., Kim, J.E., Kim, H.B., Baek, H.J., 2015. Effects of water temperature change on the hematological responses and plasma cortisol levels in growing of red spotted grouper, *Epinephelus akaara*. Dev. Reprod. 19, 19–24.
- Correa, C.F., Nobrega, R.O., Mattioni, B., Block, J.M., Fracalossi, D.M., 2017. Dietary lipid sources affect the performance of Nile tilapia at optimal and cold, suboptimal temperatures. Aquac. Nutr. 23 (5), 1016–1026.
- Corrêa, C.F., Nobrega, R.O., Block, J.M., Fracalossi, D.M., 2018. Mixes of plant oils as fish oil substitutes for Nile tilapia at optimal and cold suboptimal temperature. Aquaculture 497, 82–90. <https://doi.org/10.1016/j.aquaculture.2018.07.034>.
- Coutteau, P., Geurden, I., Camara, M.R., Bergot, P., 1997. Review on the dietary effects of phospholipids in fish and crustacean larviculture. Aquaculture 155 (1–4), 149–164.
- Coz-Rakovac, R., Strunjak-Perovic, I., Hacmanjek, M., Lipej, Z., Sostaric, B., 2005. Blood chemistry and histological properties of wild and cultured sea bass (*Dicentrarchus labrax*) in the North Adriatic Sea. Vet. Res. Commun. 29 (8), 677–687.
- Craig, S.R., Neill, W.H., Gatlin, D.M., 1995. Effects of dietary lipid and environmental salinity on growth, body composition, and cold tolerance of juvenile red drum (*Sciaenops ocellatus*). Fish. Physiol. Biochem. 14, 49–61.
- Dan, N.C., Little, D.C., 2000. The culture performance of monosex and mixed-sex newseason and overwintered fry in three strains of Nile tilapia (*Oreochromis niloticus*) in northern Vietnam. Aquaculture 184 (3–4), 221–231.
- Dogru, M.I., Dogru, A.K., Gul, M., Esrefoglu, M., Yurekli, M., Erdogan, S., Ates, B., 2008. The effect of adrenomedullin on rats exposed to lead. J. Appl. Toxicol. 28 (2), 140–146.
- Elashry, M.A., Mohammady, E.Y., Soaudy, M.R., Ali, M.M., El-Garhy, H.S., Ragaza, J.A., Hassaan, M.S., 2024. Growth, health, and immune status of Nile tilapia *Oreochromis niloticus* cultured at different stocking rates and fed algal β -carotene. Aquac. Rep. 35, 101987.
- El-faki, F.E.M., 2016. Eval. Addit. Gum. Arab. Growth Perform. Indices Mix-sex. Tilapia O. *Niloticus* Finger (Dr. Diss. Sudan Univ. Sci. Technol.).
- El-Naggar, K., Mohamed, R., El-katcha, M.I., Abdo, S.E., Soltan, M.A., 2021. Plant Ingredient diet supplemented with lecithin as fish meal and fish oil alternative affects growth performance, serum biochemical, lipid metabolism and growth-related gene expression in Nile tilapia. Aquac. Res. 52 (12), 6308–6321.
- El-Sayed, A.-F.-M., Tammam, M.S., Makled, S.O., 2021. Lecithin-containing bioemulsifier boosts growth performance, feed digestion and absorption and immune response of adult Nile tilapia (*Oreochromis niloticus*). Aquac. Nutr. 27 (3), 757–770. <https://doi.org/10.1111/anu.13221>.
- Faggio, C., Fazio, F., Marafioti, S., Arfuso, F., Piccione, G., 2015. Oral administration of Gum Arabic: effects on haematological parameters and oxidative stress markers in *Mugil cephalus*. Iran. J. Fish. Sci. 14 (1), 60–72.
- Folch, J., Lees, M., Stanley, G.S., 1957. A simple method for the isolation and purification of total lipides from animal tissues. J. Biol. Chem. 226 (1), 497–509.
- Fouda, A., El-Aziz, Abd, Mabrouk, N., 2019. Effects of arabic gum on cardiomyopathy in a rat model of type II diabetes. Al-Azhar Med. J. 48, 29–42.
- Fracalossi, D.M., Lovell, R.T., 1995. Growth and liver polar fatty acid composition of year-1 channel catfish fed various lipid sources at two water temperatures. Prog. Fish. Cult. 57, 107–113. [https://doi.org/10.1577/1548-8640\(1995\)0572.3.CO;2](https://doi.org/10.1577/1548-8640(1995)0572.3.CO;2).
- Fredenrich, A., Bayer, P., 2003. Reverse cholesterol transport, high density lipoproteins and HDL cholesterol: recent data. Diabetes Metab. 29 (3), 201–205.
- Galloway, B.J., Kieffer, J.D., 2003. The effects of an acute temperature change on the metabolic recovery from exhaustive exercise in juvenile Atlantic salmon (*Salmo salar*). Physiol. Biochem. Zool. 76 (5), 652–662.
- Garcia, D., Lima, D., da Silva, D.G.H., de Almeida, E.A., 2020. Decreased malondialdehyde levels in fish (*Astyanax altiparanae*) exposed to diesel: evidence of

- metabolism by aldehyde dehydrogenase in the liver and excretion in water. *Ecotoxicol. Environ. Saf.* 190, 110107.
- Gisbert, E., Villeneuve, L., Zambonino-Infante, J., Quazuguel, P., Cahu, C., 2005. Dietary phospholipids are more efficient than neutral lipids for long-chain polyunsaturated fatty acid supply in European sea bass *Dicentrarchus labrax* larval development. *Lipids* 40 (6), 609. <https://doi.org/10.1007/s117-5-005-1422-0>.
- Greene, D.H., Selivonchick, D.P., 1987. Lipid metabolism in fish. *Prog. Lipid Res.* 26 (1), 53–85.
- Hagar, A.F., Hazel, J.R., 1985b. The influence of thermal-acclimation on the microsomal fatty acid composition and desaturase activity of rainbow-trout liver. *Mol. Physiol.* 7 (2), 107–118.
- Haghparsat, R.J., Moghanlou, K.S., Mohseni, M., Imani, A., 2019. Effect of dietary soybean lecithin on fish performance, hemato-immunological parameters, lipid biochemistry, antioxidant status, digestive enzymes activity and intestinal histomorphometry of pre-spawning Caspian brown trout (*Salmo trutta caspius*). *Fish. Shellfish Immunol.* 91, 50–57.
- Hamre, K., Yuferra, M., Rønnestad, I., Boglione, C., Conceicao, L.E., Izquierdo, M., 2013. Fish larval nutrition and feed formulation: knowledge gaps and bottlenecks for advances in larval. *Rev. Aquacult.*
- Hassaan, M.S., Mohammady, E.Y., Soaudy, M.R., El-Garhy, H.A.S., Moustafa, M.M.A., Mohamed, S.A., Ehab, R., El-Haroun, E.R., 2019b. Effect of Silybum marianum seeds as a feed additive on growth performance, serum biochemical indices, antioxidant status, and gene expression of Nile tilapia, *Oreochromis niloticus* (L.) fingerlings. *Aquaculture* 509, 178–187.
- Hassaan, M.S., Mohammady, E.Y., Soaudy, M.R., Sabae, S.A., Mahmoud, A.M., El Haroun, E.R., 2021. Comparative study on the effect of dietary β -carotene and phycocyanin extracted from *Spirulina platensis* on immune-oxidative stress biomarkers, genes expression and intestinal enzymes, serum biochemical in Nile tilapia, *Oreochromis niloticus*. *Fish. Shellfish Immunol.* 108, 63–72.
- Hassaan, M.S., Nagar, A.G.E., Salim, H.S., Fitzsimmons, K., El-Haroun, E.R., 2019a. Nutritional mitigation of winter thermal stress in Nile tilapia by propolis-extract: associated indicators of nutritional status, physiological responses and transcriptional response of delta-9-desaturase gene. *Aquaculture* 511, 734256.
- Hassaan, M.S., Soltan, M.A., Abdel-Moez, A.M., 2015. Nutritive value of soybean meal after solid state fermentation with *Saccharomyces cerevisiae* for Nile tilapia, *Oreochromis niloticus*. *Anim. Feed Sci. Technol.* 201, 89–98.
- Henderson, R.J., Tocher, D.R., 1987. The lipid composition and biochemistry of freshwater fish. *Prog. Lipid Res.* 26 (4), 281–347.
- Hertrampf, J.W., Piedad-Pascual, F., 2000. Soya Lecithin. In *Handbook on Ingredients for Aquaculture Feeds*. Springer, pp. 383–395. https://doi.org/10.1007/978-94-011-4018-8_41.
- Hsieh, S.L., Hu, C.Y., Hsu, Y.T., Hsieh, T.J., 2007. Influence of dietary lipids on the fatty acid composition and stearyl-CoA desaturase expression in hybrid tilapia (*Oreochromis niloticus* × *O. aureus*) under cold shock. *Comp. Biochem. Physiol. Part B Biochem. Mol. Biol.* 147 (3), 438–444.
- Hurst, T.P., 2007. Causes and consequences of winter mortality in fishes. *J. Fish. Biol.* 71 (2), 315–345.
- Hyvärinen, P., Heinimaa, S., Rita, H., 2004. Effects of abrupt cold shock on stress responses and recovery in brown trout exhausted by swimming. *J. Fish. Biol.* 64 (4), 1015–1026.
- Imbabi, T.A., Habashy, W.S., Abol-Fetouh, G.M., Labib, M.M., Osman, A., Elkelish, A., Ahmed-Farid, O., 2023. Enhancing semen quality, brain neurotransmitters, and antioxidant status of rabbits under heat stress by acacia gum, vitamin C, and lycopene as dietary supplements: an in vitro and in silico study. *Ital. J. Anim. Sci.* 22 (1), 321–336.
- Inanan, B.E., Acar, Ü., Inanan, T., 2021. Effects of dietary *Ferula elaeoachyris* root powder concentrations on haematology, serum biochemical parameters, spermatozoa parameters, and oxidative status in tissues of males goldfish (*Carassius auratus*). *Aquaculture* 544, 737087. <https://doi.org/10.1016/j.aquaculture.2021.737087>.
- Jafari, F., Agh, N., Noori, F., Tokmachi, A., Gisbert, E., 2018. Effects of dietary soybean lecithin on growth performance, blood chemistry and immunity in juvenile stellate sturgeon (*Acipenser stellatus*). *Fish. Shellfish Immunol.* 80, 487–496.
- Ju, Z., Dunham, R.A., Liu, Z., 2002. Differential gene expression in the brain of channel catfish (*Ictalurus punctatus*) in response to cold acclimation. *Mol. Genet. Genom.* 268, 87–95.
- Kaushik, S., Kaur, J., 2003. Chronic cold exposure affects the antioxidant defense system in various rat tissues. *Clin. Chim. Acta* 333, 69–77.
- Kesbiç, O.S., Acar, Ü., Mohammady, E.Y., Salem, S.M.R., Ragaza, J.A., El-Haroun, E., Hassaan, M.S., 2022. The beneficial effects of citrus peel waste and its extract on fish performance and health status: a review. *Aquac. Res.* 53, 4217–4232.
- Kumar, N., Minhas, P.S., Ambasankar, K., Krishnani, K.K., Rana, R.S., 2014. Dietary lecithin potentiates thermal tolerance and cellular stress protection of milk fish (*Chanos chanos*) reared under low dose endosulfan-induced stress. *J. Therm. Biol.* 46, 40–46.
- Kyprianou, T.D., Pörtner, H.O., Anestis, A., Kostoglou, B., Feidantsis, K., Michaelidis, B., 2010. Metabolic and molecular stress responses of gilthead sea bream *Sparus aurata* during exposure to low ambient temperature: an analysis of mechanisms underlying the winter syndrome. *J. Comp. Physiol. B* 180, 1005–1018.
- Lermen, C.L., Lappe, R., Crestani, M., Vieira, V.P., Gioda, C.R., Schetinger, M.R.C., Baldisserotto, B., Moraes, G., Morsch, V.M., 2004. Effect of different temperature regimes on metabolic and blood parameters of silver catfish *Rhamdia quelen*. *Aquaculture* 239, 497–507.
- Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2- $\Delta\Delta$ CT method. *Methods* 25 (4), 402–408.
- Lortz, S., Tiedge, M., Nachtwey, T., Karlsen, A.E., Nerup, J., Lenzen, S., 2000. Protection of insulin-producing rim5f cells against cytokine-mediated toxicity through over expression of antioxidant enzymes. *Diabetes* 49, 1123–1130.
- Lucas, A., 1996. Physical concepts of bioenergetics. In: Lucas, A. (Ed.), *Bioenergetics of Aquatic Animals*, English ed. Taylor & Francis, France.
- Ma, X.Y., Qiang, J., He, J., Gabriel, N.N., Xu, P., 2015. Changes in the physiological parameters, fatty acid metabolism, and SCD activity and expression in juvenile GIFT tilapia (*Oreochromis niloticus*) reared at three different temperatures. *Fish. Physiol. Biochem.* 41, 937–950.
- Mahmoud, H.K., Reda, F.M., Alagawany, M., Farag, M.R., El-Naggar, K., 2023. The role of dietary chia seed powder in modulating cold stress-related impacts in Nile tilapia, *Oreochromis niloticus*. *Aquaculture* 567, 739246.
- Mirghani, M.E., Elnour, A.A., Kabbashi, N., Alam, M.Z., Musa, K.H., Abdullah, A., 2018. Determination of antioxidant activity of gum arabic: an exudation from two different locations. *Sci. Asia* 44, 179–186.
- Moawad, D.A., El-Garhy, H.A., Refaat, M.H., Hassaan, M.S., 2024. Protecting deleterious effects of high stocking density of Nile tilapia, *Oreochromis niloticus* using dietary microalgae *Golenkinia longispicula*: growth, immune-oxidative markers and associated gene expression. *Aquaculture* 581, 740430.
- Mohammady, E.Y., Aboseif, A.M., Soaudy, M.R., Ramadan, E.A., Hassaan, M.S., 2023. Appraisal of fermented wheat bran by *Saccharomyces cerevisiae* on growth, feed utilization, blood indices, intestinal and liver histology of Nile tilapia, *Oreochromis niloticus*. *Aquaculture*, 739755.
- Mohammady, E.Y., Elashry, M.A., Ibrahim, M.S., Elarian, M., Salem, S.M.R., El-Haroun, E.R., Mohamed, S., Hassaan, M.S., 2024. Nano iron versus bulk iron forms as functional feed additives: growth, body indices, hematological assay, plasma metabolites, immune, anti-oxidative ability, and intestinal morphometric measurements of Nile tilapia, *Oreochromis niloticus*. *Biol. Trace Elem. Res.* 202, 787–799.
- Moin, V.M., 1986. A simple and specific method for determining glutathione peroxidase activity in erythrocytes. *Lab. Invest.* 55, 724–727.
- Mommsen, T.P., Vijayan, M.M., Moon, T.W., 1999. Cortisol in teleosts: dynamics, mechanisms of action, and metabolic regulation. *Rev. Fish. Biol. Fish.* 9 (3), 211–268.
- Morais, S., Conceicao, L., Rønnestad, I., Koven, W., Cahu, C., Infante, J.Z., Dinis, M., 2007. Dietary neutral lipid level and source in marine fish larvae: effects on digestive physiology and food intake. *Aquaculture* 268 (1–4), 106–122. <https://doi.org/10.1016/j.aquaculture.2007.04.033>.
- Musa, K.H., Abdullah, A., Kuswandi, B., Hidayat, M.A., 2013. A novel high throughput method based on the DPPH dry reagent array for determination of antioxidant activity. *Food Chem.* 141, 4102–4106.
- Naiel, M.A., Abd El-Hameed, S.A., Arisha, A.H., Negm, S.S., 2022. Gum Arabic-enriched diet modulates growth, antioxidant defenses, innate immune response, intestinal microbiota and immune related genes expression in tilapia fish. *Aquaculture* 556, 738249.
- Naiel, M.A., Ismael, N.E., Negm, S.S., Ayyat, M.S., Al-Sagheer, A.A., 2020. Rosemary leaf powder-supplemented diet enhances performance, antioxidant properties, immune status, and resistance against bacterial diseases in Nile Tilapia (*Oreochromis niloticus*). *Aquaculture* 526, 735370.
- Navruz, F.Z., Acar, Ü., Yilmaz, S., Kesbiç, O.S., 2023. Dose-dependent stress response of esfenvalerate insecticide on common carp (*Cyprinus carpio*): evaluating blood parameters and gene expression. *Comp. Biochem. Physiol. Part C Toxicol. Pharmacol.* 272, 109711. <https://doi.org/10.1016/j.cbpc.2023.109711>.
- Nobrega, R.O., Batista, R.O., Corrêa, C.F., Mattioni, B., Filer, K., Pettigrew, J.E., Fracalossi, D.M., 2019. Dietary supplementation of *Aurantiochytrium* sp. meal, a docosahexaenoic acid source, promotes growth of Nile tilapia at a suboptimal low temperature. *Aquaculture* 507, 500–509. <https://doi.org/10.1016/j.aquaculture.2019.04.030>.
- Nobrega, R.O., Corrêa, C.F., Mattioni, B., Fracalossi, D.M., 2017. Dietary α -linolenic for juvenile Nile tilapia at cold suboptimal temperature. *Aquaculture* 471, 66–71. <https://doi.org/10.1016/j.aquaculture.2016.12.026>.
- Osman, A.G., Koutb, M., Sayed, A.E.D.H., 2010. Use of hematological parameters to assess the efficiency of quince (*Cydonia oblonga* Miller) leaf extract in alleviation of the effect of ultraviolet-a radiation on African catfish *Clarias gariepinus* (Burchell, 1822). *J. Photochem. Photobiol. B Biol.* 99 (1), 1–8.
- Pacheco, M., Santos, M.A., 2001. Biotransformation, endocrine, and genetic responses of *Anguilla anguilla* L. to petroleum distillate products and environmentally contaminated waters. *Ecotoxicol. Environ. Saf.* 49, 64–75.
- Panase, P., Saenphet, S., Saenphet, K., 2018. Biochemical and physiological responses of Nile tilapia, *Oreochromis niloticus* Lin subjected to cold shock of water temperature. *Aquac. Rep.* 11, 17–23.
- Peskin, A.V., Winterbourn, C.C., 2000. A microtiter plate assay for superoxide dismutase using a water-soluble tetrazolium salt (WST-1). *Clin. Chim. Acta* 293 (1–2), 157–166.
- Prieto, P., Pineda, M., Aguilar, M., 1999. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. *Anal. Biochem.* 269 (2), 337–341.
- Radwan, I., 2008. Tilapia aquaculture in the Nile Delta 1990–2008. *Proceedings of 8th International Symposium on Tilapia in Aquaculture, Cairo, Egypt* 12–14. 10.1111/j.1753-5131.2012.01086.x.
- Refaey, M.M., Mehri, A.I., Zenhom, O.A., Mansour, A.T., 2022. Effect of fatty acids manipulation on survival and physiological response of hybrid red tilapia under chronic cold stress. *Aquaculture* 561, 738663.
- Reitman, S., Frankel, S., 1957. A colorimetric method for the determination of serum glutamic oxalacetic and glutamic pyruvic transaminases. *Am. J. Clin. Pathol.* 28 (1), 56–63.

- Rothbard, S., Peretz, Y., 2002. Tilapia Culture in NEGEV, the Israeli Desert. In: Guerrero, R.D., R. Guerrero-del Castillo (Eds.), *Tilapia Farming in the 21st Century*, pp. 60–65.
- Rudneva, I.I., Kuzminova, N.S., Skuratovskaya, E.N., 2010. Glutathione-S-transferase activity in tissues of Black Sea fish species. *Asian J. Exp. Biol. Sci.* 1, 141–150.
- Ruyter, B., Røjo, C., Grisdale-Helland, B., Rosenlund, G., Obach, A., Thomassen, M.S., 2003. Influence of temperature and high dietary linoleic acid content on esterification, elongation, and desaturation of PUFA in Atlantic salmon hepatocytes. *Lipids* 38 (8), 833–840.
- Saleh, R.S., Mohammady, E.Y., El-Haroun, E., Hassaan, M.S., 2022. Dietary dried periphyton can improve growth, digestive enzyme, serum biochemical, antioxidant response and intestinal morphometric of Nile tilapia. *Aquac. Res.* 53 (18), 6463–6477.
- Sandamali, W.G., Kwon, H., Lim, C., Yang, H., Lee, J., 2021. The possible role of catalase in innate immunity and diminution of cellular oxidative stress: insights into its molecular characteristics, antioxidant activity, DNA protection, and transcriptional regulation in response to immune stimuli in yellowtail clownfish (*Amphiprion clarkii*). *Fish. Shellfish Immunol.* 113, 106–117.
- Šegvić-Bubić, T., Boban, J., Grubišić, L., Trumbić, Ž., Radman, M., Perčić, M., Čož-Rakovac, R., 2013. Effects of propolis enriched diet on growth performance and plasma biochemical parameters of juvenile European sea bass (*Dicentrarchus labrax* L.) under acute low-temperature stress. *Aquac. Nutr.* 19 (6), 877–885.
- Seiliez, I., Bruant, J.S., Zambonino Infante, J.L., Kaushik, S., Bergot, P., 2006. Effect of dietary phospholipid level on the development of gilthead sea bream (*Sparus aurata*) larvae fed a compound diet. *Aquac. Nutr.* 12, 372–378.
- Shang, H.S., Shih, Y.L., Chen, C.P., Lee, M.H., Lu, H.F., Chou, P.Y., Liao, N.C., Chen, Y.L., Hsueh, S.C., Chung, J.G., 2018. Laminarin Promotes Immune Responses and Normalizes Glutamic Oxaloacetic Transaminase and Glutamic Pyruvic Transaminase Levels in Leukemic Mice in Vivo. *Vivo* 32 (4), 783–790. <https://doi.org/10.21873/in vivo.11308>.
- Shi, G.C., Dong, X.H., Chen, G., Tan, B.P., Yang, Q.H., Chi, S.Y., Liu, H.Y., 2015. Physiological responses and HSP70 mRNA expression of GIFT strain of Nile tilapia (*Oreochromis niloticus*) under cold stress. *Aquac. Res.* 46, 658–668.
- Siddik, M.A.B., Nahar, A., Ahamed, F., Hossain, Md.Y., 2014. Over-wintering growth performance of mixed-sex and mono-sex Nile tilapia *Oreochromis niloticus* in northeastern Bangladesh. *Croat. J. Fish.* 72, 70–76.
- Sivaramakrishnan, T., Ambasankar, K., Kumaraguru Vasagam, K.P., Syama Dayal, J., Sandeep, K.P., Bera, A., Magesh, M., Kailasam, M., Vijayan, K.K., 2021. Effect of dietary soy lecithin inclusion levels on growth, feed utilization, fatty acid profile, deformity and survival of milkfish (*Chanos chanos*) larvae. *Aquac. Res.* 52, 5366–5374. <https://doi.org/10.1111/are.15406>.
- Soaudy, M.R., Mohammady, E.Y., Elashry, M.A., Ali, M.M., Ahmed, N.M., Hegab, M.H., El-Garhy, H.A., El-Haroun, E.R., Hassaan, M.S., 2021. Possibility mitigation of cold stress in Nile tilapia under biofloc system by dietary propylene glycol: performance feeding status, immune, physiological responses and transcriptional response of delta-9-desaturase gene. *Aquaculture* 538, 736519.
- Soltan, M.A., Abdella, M.M., El-Sayaad, G.A., Abu-El Wafa, M.H., 2015. Effect of some over-wintering regimes on survival and growth performance of Nile tilapia *Oreochromis niloticus*. *Biol. Chem. Res.* 2, 362–374.
- Sun, Z., Tan, X., Liu, Q., Ye, H., Zou, C., Xu, M., Ye, C., 2019. Physiological, immune responses and liver lipid metabolism of orange-spotted grouper (*Epinephelus coioides*) under cold stress. *Aquaculture* 498, 545–555.
- Suski, C.D., Killen, S.S., Kieffer, J.D., Tufts, B.L., 2006. The influence of environmental temperature and oxygen concentration on the recovery of largemouth bass from exercise: implications for live-release angling tournaments. *J. Fish. Biol.* 68 (1), 120–136.
- Taalab, H.A., Mohammady, E.Y., Hassan, T.M., Abdella, M.M., Hassaan, M.S., 2022. β -Carotene of *Arthrospira platensis* versus vitamin C and vitamin E as a feed supplement: effects on growth, haemato-biochemical, immune-oxidative stress and related gene expression of Nile tilapia fingerlings. *Aquac. Res.* 53 (13), 4832–4846.
- Tandon, R.S., Joshi, B.D., 1974. Effect of trypanosome infection on blood glucose levels of some fresh water teleosts. *J. Inland Fish. Soc. India* 6, 81–82.
- Tocher, D.R., Bell, J.G., Sargent, J.R., 1996. Induction of Delta 9-fatty acyl desaturation in rainbow trout, *Oncorhynchus mykiss*, liver by dietary manipulation. *Comp. Biochem. Physiol. B Biochem. Mol. Biol.* 113 (1), 205–212.
- Tocher, D.R., Bendiksen, E.Å., Campbell, P.J., Bell, J.G., 2008. The role of phospholipids in nutrition and metabolism of teleost fish. *Aquaculture* 280 (1), 21–34. <https://doi.org/10.1016/j.aquaculture.2008.04.034>.
- Trushenski, J., Schwarz, M., Pessoa, W.V.N., Mulligan, B., Crouse, C., Gause, B., et al., 2013. Amending reduced fish-meal feeds with marine lecithin, but not soy lecithin, improves the growth of juvenile cobia and may attenuate heightened responses to stress challenge. *J. Anim. Physiol. Anim. Nutr.* 97, 170–180. <https://doi.org/10.1111/j.1439-0396.2011.01255.x>.
- Turchini, G.M., Torstensen, B.E., Ng, W.K., 2009. Fish oil replacement in finfish nutrition. *Rev. Aquac.* 1, 10–57. <https://doi.org/10.1111/j.1753-5131.2008.01001.x>.
- Tüzgen, S., Kaynar, M.Y., Güner, A., Gümüştas, K., Belce, A., Etus, V., Ozyurt, E., 1998. The effect of epidural cooling on lipid peroxidation after experimental spinal cord injury. *Spinal Cord* 36, 654–657.
- Tyagi, A., Ali, D.S., Barat, D.A., 2013. Impact of exposure of cold temperature on hematological parameters and plasma electrolytes of cold water fish *Shizothorax richardsonii*. *J. Inf. Knowl. Res. Biomed. Eng.* 2 (2), 38–41.
- Vanlandeghem, M.M., Wahl, D.H., Suski, C.D., 2010. Physiological responses of largemouth bass to acute temperature and oxygen stressors. *Fish. Manag. Ecol.* 17 (5), 414–425.
- Vijayan, M.M., Pereira, C., Graul, E.G., Lwama, G.K., 1997. Metabolic responses to confinement stress in tilapia: the role of cortisol. *Comp. Biochem. Physiol.* 116C, 89–95.
- Weber, T.E., Bosworth, B.G., 2005. Effects of 28 day exposure to cold temperature or feed restriction on growth, body composition, and expression of genes related to muscle growth and metabolism in channel catfish. *Aquaculture* 246 (1–4), 483–492.
- Wendelaar Bonga, S.E., 1997. The stress response in fish. *Physiol. Rev.* 77 (3), 591–625.
- Weyh, C., Krüger, K., Peeling, P., Castell, L., 2022. The role of minerals in the optimal functioning of the immune system. *Nutrients* 14 (3), 644.
- Wu, D., Liu, Z., Yu, P., Huang, Y., Cai, M., Zhang, M., Zhao, Y., 2020. Cold stress regulates lipid metabolism via AMPK signalling in *Cherax quadricarinatus*. *J. Therm. Biol.* 102693.
- Wu, F., Yang, C., Wen, H., Chen Zhang, C., Jiang, M., Liu, W., Tian, J., Yu, L., Xing Lu, X., 2019. Improving low-temperature stress tolerance of tilapia, *Oreochromis niloticus*: a functional analysis of *Astragalinus membranaceus*. *J. World Aquacult. Soc.* 50, 749–762.
- Yang, J.L., Chen, H.C., 2003. Effects of gallium on common carp (*Cyprinus carpio*): acute test, serum biochemistry, and erythrocyte morphology. *Chemosphere* 53 (8), 877–882.
- Yang, S., Zhang, C., Xu, W., Li, D., Feng, Y., Wu, J., Luo, W., Du, X., Du, Z., Huang, X., 2022. Heat stress decreases intestinal physiological function and facilitates the proliferation of harmful intestinal microbiota in sturgeons. *Front. Microbiol.* 13, 755369.
- Yao, J., Wang, J.Y., Liu, L., Li, Y.X., Xun, A.Y., Zeng, W.S., Jia, C.H., Wei, X.X., Feng, J.L., Zhao, L., Wang, L.S., 2010. Anti-oxidant effects of resveratrol on mice with DSS-induced ulcerative colitis. *Arch. Med. Res.* 41 (4), 288–294.
- Ye, C.X., Wan, F., Sun, Z.Z., Cheng, C.H., Ling, R.Z., Fan, L.F., Wang, A.L., 2015. Effect of phosphorus supplementation on cell viability, anti-oxidative capacity and comparative proteomic profiles of puffer fish (*Takifugu obscurus*) under low temperature stress. *Aquaculture* 452, 200–208.
- Yokoyama, S., Koshio, S., Takakura, N., Oshida, K., Ishikawa, M., Gallardo-Cigarroa, F.J., Teshima, S.I., 2005. Dietary bovine lactoferrin enhances tolerance to high temperature stress in Japanese flounder *Paralichthys olivaceus*. *Aquaculture* 249 (1–4), 367–373.
- Zerai, D.B., Fitzsimmons, K.M., Collier, R.J., 2010. Transcriptional response of delta-9-desaturase gene to acute and chronic cold stress in Nile tilapia, *Oreochromis niloticus*. *J. World Aquacult. Soc.* 41 (5), 800–806.
- Zhai, S.W., Lu, J.J., Chen, X.H., 2014. Effects of dietary grape seed proanthocyanidins on growth performance, some serum biochemical parameters and body composition of tilapia (*Oreochromis niloticus*) fingerlings. *Ital. J. Anim. Sci.* 13, 3357.
- Zikić, R., Stajin, A.S., Pavlović, S.Z., Ognjanović, B.I., Saičić, Z., 2001. Activities of superoxide dismutase and catalase in erythrocytes and plasma transaminases of goldfish (*Carassius auratus gibelio* Bloch.) exposed to cadmium. *Physiol. Res.* 50, 105–111.