

Induction of Resistance in Watermelon Plants against Fusarium Wilt using Chemical Inducers and Compost under Greenhouse Conditions

A.M.M. Mahdy*; M.H. Abd-El-Mageed*;
Faten M. Abd-El-Latif*; M.M.M. Diab**
and Nehal M. Saied**

* Plant Pathol. Dept., Fac. Agric., Benha Univ., Egypt.

** Plant Pathol. Dep., N.R.C., Egypt.

Fusarium wilt of watermelon caused by *Fusarium oxysporum* f.sp. *niveum* is the most severe disease attacking watermelon plants. Induced resistance in watermelon plants using chitosan, salicylic acid and compost was evaluated. All tested isolates of *F. oxysporum* f.sp. *niveum* were able to attack watermelon plants causing damping-off symptoms. The most aggressive isolate was isolated from Nobariya (cv. Giza 1). It caused 71.9, 46.9 and 51.6% for cvs. Gorma, Giza1 and Aswan hybrid, respectively, 30 days after sowing. Laboratory experiment results revealed that all tested concentrations of chitosan and salicylic acid significantly reduced the disease incidence under greenhouse conditions. Moreover, animal and plant compost at all concentrations significantly reduced the watermelon wilt incidence. Plant compost at 10.0 g/kg soil, chitosan at 8.0 g/kg soil and salicylic acid at 2.0 g/kg soil were tested alone or in combination to study their effect on watermelon wilt incidence and on enzyme activities of watermelon transplants. Results showed that the highest reduction in disease incidence was recorded with combined treatments between plant compost and chitosan at concentration of 8.0 g/kg soil or salicylic acid at 2.0 g/kg soil which reduced the wilt incidence by 92.9, 88.5 and 86.6% for Aswan hybrid, cvs. Gorma and Giza1, respectively. As for enzyme activities, results revealed that the highest increase was recorded with combined treatments between plant compost and chitosan at concentration of 8 g/kg soil or salicylic acid at 2.0 g/kg soil which increased the peroxidase, chitinase and β -1,3-glucanase activities more than 175.0, 200.0 and 201.0%, respectively.

Keywords: β -1,3-glucanase, chitinase, chitosan, compost, Fusarium wilt, peroxidase, salicylic acid and watermelon.

Watermelon (*Citrullus lanatus* (Trunb.) Matsum and Nakai) is a widely cultivated vegetable crop that consumed globally as fruits. The fungus *Fusarium oxysporum* Schleicher: Fr. f.sp. *niveum* (E.F. Smith) W.C. Snyder and H.N. Hans is the causal of watermelon Fusarium wilt (Booth, 1971) and found to be worldwide in tropical and subtropical regions. Watermelon wilt disease appear during the growing season at different stages of plant growth from seedling to mature stages and may happen earlier to cause pre-emergence damping-off (Sheng *et al.*, 2009 and Lu *et al.*, 2014). Also, Booth (1971) reported that *F. oxysporum* f.sp. *niveum* causes damping-off, stunting of watermelon seedlings and wilt of older plants. An

investigation of this disease is considered important especially in view of its wide prevalence in Egypt particularly in sandy soils where watermelon is an important crop.

Recently fungicide alternatives are promising methods for controlling plant diseases. Induced resistance is accomplished by the inoculation of plant with an avirulent or non-pathogenic isolates prior to or concomitant with a challenge inoculation with a pathogen. That is called biotic inducers, while chemical inducers include natural or synthetic chemicals, *i.e.* ethephon, acetylsalicylic acid, salicylic acid and chitosan (Suprakash and Chatterjee, 2012; Abd-El-Kareem and Abd-El-Latif, 2012 and Abd-El-Kareem *et al.*, 2013a).

Chitosan applied as seed or soil treatments was reported to control *Fusarium* wilts in many plant species (Badawy *et al.*, 2005). Chitosan induces host defence responses in both monocotyledons and dicotyledons (Elwagia and Algam, 2014 and Mishra *et al.*, 2014). Moreover, Salicylic acid ($C_7H_6O_3$) is an important signalling molecule involved in both locally and systemically induced disease resistance responses. The ability to accumulate salicylic acid has been shown to be essential for systemic acquired resistance and reactions to abiotic stress in plants (Morse *et al.*, 2007; Zawoznik *et al.*, 2007 and Fawzy, 2013).

On the other hand, utilization of composts to minimize organic waste pollution and to reduce the addition of chemical fertilizers and fungicides in crop production is a promising strategy for both the present and the future. Furthermore, many soilborne pathogens can be reduced by application of composts made of different raw materials (Trillas *et al.*, 2006; Abd-El-Kareem *et al.*, 2013b and Zhao *et al.*, 2014).

The present research was designed to study and evaluate the effect of chitosan, salicylic acid and compost, alone or in combination, against watermelon wilt incidence under greenhouse conditions.

Materials and Methods

Isolation, identification and Pathogenicity test:

Isolation trails were carried out from watermelon plants cvs. Gorma, Giza 1 and Aswan hybrid showing wilt symptoms collected from different locations, *i.e.* Kafr El-Shikh (Baltem & Kafr El-Shikh), Ismailiya (Abo-Soyer & Salhiya) and Behira (Nobariya & Badr City). The obtained isolates were purified and identified according to Gilman (1957) and Booth (1971). Three identified isolates of *Fusarium oxysporum* isolated from the three watermelon cultivars of each location were tested for their pathogenic ability on watermelon plants under greenhouse conditions. Experiment was carried out in Plant Pathology greenhouse at the National Research Centre, Dokki, Egypt. Inocula of *F. oxysporum* isolates were prepared by culturing each of the isolate in 50 mL potato dextrose broth (PDB) medium in 250 mL Erlenmeyer flasks for 15 days at ($25\pm 2^\circ\text{C}$). Inoculum of each *F. oxysporum* isolate was prepared from the growing upper solid layers which were blended in sterilized water. Colonies forming units (cfu) were adjusted to 10^6 cfu/mL using

haemocytometer slide. Soil infestation was carried out at the rate of 50 mL (10^6 cfu/mL)/kg soil (Elad and Baker, 1985). Plastic pots (30-cm-diam., 5.0 kg soil) containing sterilized sandy-loamy soil autoclaved at 120°C for 1h were artificially infested individually with the inoculum of the desired isolate at the rate of 50 mL (10^6 cfu/mL)/kg soil (Elad and Baker, 1985). Eight pots were used as replicates for each isolate as well as check treatment (uninfested soil). Disinfested watermelon seeds of cvs. Giza1, Gorma and Aswan hybrid were sown at the rate of 8 seeds/pot. Disease incidence was estimated as percentage of dead plants 15 and 30 days after sowing.

Assessment of dead plants:

Fusarium damping-off was measured as percentage of dead plants at 15 and 30 days after sowing (Booth, 1971) as follow:

$$\text{Percentage of dead plants (\%)} = \frac{\text{Number of dead plants}}{\text{Total number of planted seeds}} \times 100$$

Host range of F. oxysporum isolated from watermelon:

The highest aggressive isolate of *F. oxysporum* causing watermelon wilt incidence was chosen for testing its ability to induce wilt disease on various plant species belonging to different families. This experiment was carried out under greenhouse conditions. Ten plant species belonging to families *Cucurbitaceae*, *Solanaceae* and *Leguminosae* were tested. Three watermelon cultivars, i.e. Giza 1, Gorma and Aswan hybrid, in addition to cucumber (*Cucumis sativus* L.) cv. Beit Alpha, Muskmelon (*Cucumis melo* L.) cv. Honeydew, Squash (*Cucurbita pepo* L.) cv. Eskandarany, tomato (*Solanum lycopersicum* L.) cv. Castel Rock, pepper (*Piper nigrum* L.) cv. California, bean (*Phaseolus vulgaris* L.) cv. Giza 3 and pea (*Pisum sativum* L.) cv. Master were tested. Inoculum preparation, soil infestation and assessment of disease incidence were carried out as abovementioned, 18, 30 and 45 days after sowing.

In vitro evaluation of the inhibitory effect of chitosan and salicylic acid on the linear growth and spore germination of F. oxysporum f.sp. niveum:

The inhibitory effect of chitosan and salicylic acid (Sigma Company) on the linear growth and spore germination of the *F. oxysporum* f.sp. *niveum* was evaluated under laboratory conditions. Five concentrations of chitosan solutions, i.e. 0, 2, 4, 6 and 8 g/l and five concentrations of salicylic acid, i.e. 0, 0.5, 1.0, 1.5 and 2.0 g/l (these concentrations are equivalent to be as 0.0, 3.6, 7.2, 10.9 and 14.5 mM) were tested. As for linear growth test, chitosan and salicylic acid solutions were added individually to conical flasks containing sterilized PDA medium before its solidification and mixed gently then transferred in sterilized Petri plates (9-cm diam.). Plates were individually inoculated with equal disks (6-mm-diam.) taken from 7 days old cultures of *F. oxysporum* f.sp. *niveum*, then incubated at 25±2°C. Linear growth of the fungus was measured, when the control plates reached full growth and the average growth diameter was calculated. Each treatment was represented by 5 plates as replicates. As for spore germination test, spore suspension was prepared by culturing *F. oxysporum* f. sp. *niveum* in Petri plates containing PDA

medium for 20 days at $25\pm 2^{\circ}\text{C}$. Colony forming units (cfu) containing hyphal fragments, microconidia, macroconidia and chlamydo spores were released in sterilized water using a needle then adjusted to 10^6 cfu/mL using haemocytometer slide. One mL of the suspension was transferred to test tube containing sterilized PDB (broth) medium treated with previous concentrations of chitosan or salicylic acid. Test tubes were incubated for 24 h at $25\pm 2^{\circ}\text{C}$. One mL of treated spore suspension (cfu) was examined microscopically and the percent of spore germination was calculated.

Disease control experiments:

The efficacy of the tested chemicals and compost used for controlling damping-off incidence of watermelon was carried out in pot experiment.

Watermelon seeds:

Watermelon seeds (cv. Giza1) were obtained from the Dept. of Vegetable Crop Res., Agric. Res. Centre, Giza. While, Aswan hybrid was obtained from Sakata Company, Japan and cv. Gorma from commercial markets in Egypt.

Evaluation of chitosan and salicylic acid on watermelon damping-off incidence:

Seed treatment with chitosan or salicylic acid:

Watermelon seeds cvs. Giza1, Gorma and Aswan hybrid were soaked individually for 24 hours in each solution of chitosan at concentrations of 0, 2, 4, 6 and 8 g/l or salicylic acid solutions at concentrations of 0.0, 0.5, 1.0, 1.5 and 2.0 g/l then kept between two heavy layers of cottony cheesecloth saturated with tap water for 24 h at 30°C until the beginning of germination. Seeds soaked in tap water served as control.

Soil treatment with chitosan or salicylic acid:

Artificially potted infested soil was treated by chitosan at the rate 0.0, 2.0, 4.0, 6.0 and 8.0 g/kg of soil, while, salicylic acid was used at the rate of 0.0, 0.5, 1.0, 1.5 and 2.0 g/kg soil.

Effect of soil amendment with compost for controlling Fusarium damping-off:

This study was carried-out in pots contained artificially infested soil with *F. oxysporum niveum*. Two types of composts (obtained from El-Nile Company, Giza, Egypt), *i.e.* animal and plant composts at concentrations of 0.0, 5.0 and 10.0 g/kg soil were used as soil drench 15 days before sowing to evaluate their effects on Fusarium damping-off incidence of watermelon. Disinfected watermelon seeds were sown at the rate of 8 seeds/pot and 8 pots were used for every treatment. Disease incidence was assessed as mentioned before.

Efficacy of chitosan or salicylic acid alone or in combination with compost on Fusarium wilt incidence of watermelon transplants under greenhouse conditions:

Chitosan at 8 g/kg soil or salicylic acid at 2 g/kg were applied as seed bed (treated peat-moss in trays 84 eyes before seed sowing). Disinfected watermelon seeds, cvs. Giza 1, Gorma and Aswan hybrid were sown in treated seed bed. Watermelon transplants (25-day-old) were in transplanted in plastic pots (30-cm-diam.) containing infested soil with the pathogenic planted fungus and treated with

plant compost at the rate of 10.0 g/kg soil. Transplants were transplanted at the rate of 4 transplants/pot and 8 replicates were used.

Disease assessment:

Number of wilted plants at 30 days after transplanting was recorded and the percentage of disease incidence was calculated, by using the disease index scale: 1, apparently healthy plants ; 2, slight chlorosis of lower ; slight wilt of plants ; 3 necrosis, following of lower leaves, yellow areas on upper leaves ; 4 dead plant. (Tziros, *et al.*, 2007).

Determination of enzymes activities:

Watermelon plants cvs. Giza1, Gorma and Aswan hybrid (20-day after transplanting) representing the different treatments grown under greenhouse conditions were used to determine the activity of some enzymes related to plant resistance, *i.e.* peroxidase, chitinase and β -1,3-glucanase.

Extraction of enzymes:

Plant roots (g) were homogenized with 0.1 M sodium phosphate buffer (pH 7.1) (Goldschmidt *et al.*, 1968) at the rate of 1/3 w/v. The homogenate was centrifuged at 3000 rpm for 15 min. The supernatant was used to determine enzyme activities.

Peroxidase assay:

Peroxidase activity was measured by incubation 0.1 mL of enzyme extract with 4mL of guaiacol solution for one minute at 25°C and absorbance at 470 nm was determined. The guaiacol solution consisted of 3 mL of 0.05 M K. phosphate, pH 7, 0.5 mL of 2% guaiacol and 0.5 mL of 0.3% H₂O₂ (Abeles *et al.*, 1971). Peroxidase activity was expressed as the increase in absorbance at 470 nm/g fresh weight/1 min.

Chitinase assay:

The substrate colloidal chitin was prepared from chitin powder according to the method described by Ried and Ogryd-Ziak (1981). Determination the chitinase activity was carried out according to the method of Monreal and Reese (1969), 1 mL of 1% colloidal chitin in 0.05 M citrate phosphate buffer (pH 6.6) in test tubes, 1 mL of enzyme extract was added and mixed by shaking. Tubes were kept in a water bath at 37°C for 60 min, then cooled and centrifuged before assaying. Reducing sugar was determined in 1 mL of the supernatant by dinitrosalicylic acid (DNS). The reaction was stopped by heating the tubes for 5 min at 100°C. The tubes were cooled and 3 mL distilled water were added before assay. Optical density was determined at 540 nm. Chitinase activity was expressed as mM N-acetylglucosamine equivalent released/g fresh weight tissue/60 min.

β -1,3-glucanase assay:

The method of Abeles and Forrence (1970) was used to determine β -1,3-glucanase activity. Laminarin was used as substrate and dinitrosalicylic acid as reagent to measure reducing sugars. The method was carried out as 0.5 mL of enzyme extract was added to 0.5 mL of 0.05 M of potassium acetate buffer (pH 5) containing 2% laminarin. The mixture was incubated at 40°C for 60 min. The reaction was stopped by adding 1 mL of dinitrosalicylic acid reagent and heating the tubes for 5 min at 100°C. The tubes were cooled and 3 mL distilled water were

added before assay. The optical density was read at 500 nm. β -1,3-glucanase activity was expressed as mM glucose equivalent released/g fresh weight tissues/60 min.

Statistical analysis:

Tukey test for multiple comparisons among means was utilized (Neler *et al.*, 1985).

Results

Isolation, identification and pathogenicity test:

Isolation trails resulted in different 54 fungal isolates. The obtained isolates were screened microscopically for genus *Fusarium*. Eighteen isolates were found to be *F. oxysporum* according to Gilman (1957) and (Booth 1971). Thereafter, three identified isolates of *F. oxysporum* isolated from the three watermelon cultivars of each location were tested for their pathogenic ability to induce damping-off and/or wilt diseases to watermelon plants under greenhouse conditions.

Results in Table (1) indicate that the all isolates of *F. oxysporum* were able to attack watermelon plants causing damping off symptoms.

Table 1. Pathogenic ability of three isolates of *F. oxysporum* to induce damping-off of watermelon plants 15 and 30 days after sowing

Governorate	Isolate	Dead plants (%)					
		Watermelon cultivar					
		Gorma		Giza 1		Aswan hybrid	
		Days after sowing					
		15	30	15	30	15	30
Behira	Nobariya Aswan	18.5 b*	43.8 cd	10.9 d	32.8 c	10.9 c	32.8
	Nobariya Giza 1	28.1 a	71.9 a	18.5 a	46.9 a	20.3 a	51.6 a
	Nobariya Gorma	17.2 b	45.3 cd	15.6 bc	32.8 c	14.1 b	32.8 c
	Bader Aswan	21.9 b	60.9 b	14.1 bc	42.2 b	12.5 b	40.6 b
	Bader Giza 1	17.2b	43.8 cd	14.1 bc	29.7 c	14.1b	32.8 cd
	Bader Gorma	21.9 b	43.8 cd	12.5cd	32.8 c	12.5 b	32.8 cd
Ismailiya	Abo-Soyer Aswan	17.2b	39.1e	15.6 bc	34.4 c	12.5 bc	31.3 cd
	Abo-Soyer Giza 1	14.1 c	31.3 f	10.9d	32.8 c	12.5 b	32.8 c
	Abo-Soyer Gorma	17.2 b	39.1 e	12.5 cd	32.8 c	14.1 b	32.8 c
	Salhiya Aswan	9.4 d	31.3f	10.9d	14.1 e	10.9 c	18.8 f
	Salhiya Giza 1	12.5 d	40.6 de	10.9d	31.3 c	12.5 bc	23.4 f
Kafr El-Sheikh	Salhiya Gorma	12.5 d	39.1 e	12.5 cd	23.4c	14.1 b	32.8 c
	Baltem Aswan	17.2 b	42.2 de	12.5 cd	31.3 c	14.1 b	31.3 cd
	Baltem Giza 1	9.4 ed	31.3 f	9.4 d	23.4 d	14.1b	42.2 b
	Baltem Gorma	14.1 c	39.1 e	9.4 d	31.3 c	12.5 bc	32.8 c
	Kafr El-Sheikh Aswan	21.9 b	53.1 c	15.6 d	32.8 c	14.1b	28.1 d
	Kafr El-Sheikh Giza 1	21.9 b	39.1 f	12.5c	23.4 d	10.9 c	21.9 f
	Kafr El-Sheikh Gorma	17.2 b	39.1 e	12.5 c	32.8 c	12.5 bc	21.9 f
	Control	3.1 f	4.7 g	3.1 e	3.1 f	3.1	3.1 e

* Figures with the same letter are not significantly different (P= 0.05).

The tested isolates significantly varied in their ability to cause damping-off infection of watermelon. The most aggressive isolate was isolated from Nobariya (cv. Giza 1) which caused damping off 28.1, 18.5 and 20.3% after 15 days from sowing for cvs. Gorma, Giza1 and Aswan hybrid, respectively. Moreover, the percentage of dead plants reached 71.9, 46.9 and 51.6% for cvs. Gorma, Giza1 and Aswan hybrid, respectively, 30 days after sowing. Other isolates caused disease infection in range 9.4-21.9 and 18.2-61.0% after 15 and 30 days from sowing, respectively. The highest infection was recorded in case of cv. Gorma followed by the other two tested watermelon cultivars. The most aggressive isolate (Nobariya - Giza 1) was used in the further studies.

Host range of F. oxysporum f.sp. niveum:

The host range test of *F. oxysporum f.sp. niveum* was carried out under greenhouse condition on different hosts as mentioned before. Results in Table (2) show that among the ten tested plants, only watermelon cultivars were susceptible to infection with the *F. oxysporum f. sp. niveum*. Cultivar Gorma was more susceptible to infection, it recorded 26.7, 73.4 and 84.4% as damping off and wilted plants 15, 30 and 45 days after sowing, respectively. Meanwhile, the tested plant cultivars of cucumber, muskmelon, squash, tomato, pepper, bean and pea plants could not be affected by the tested fungal isolate and no disease symptoms were seen. Therefore, it could be concluded that the tested fungal isolate is *F. oxysporum f.sp. niveum* (E.F. Smith) W.C. Snyder and H.N. Hans (Booth, 1971).

Table 2. Evaluation of host range of *F. oxysporum* against different plant species

Tested plant species	Fusarium wilt disease incidence (%)		
	Days after sowing		
	Damping off		Wilt
	15	30	45
Watermelon (Giza 1)	18.8 b*	60.9 b	73.4 b
Watermelon (Gorma)	26.7 a	73.4 a	84.4 a
Watermelon (Aswan)	17.2 b	62.5 b	73.4 b
Cucumber (Beit Alpha)	0.0 c	0.0 c	0.0 c
Muskmelon (Honeydew)	0.0 c	0.0 c	0.0 c
Squash (Eskandarany)	0.0 c	0.0 c	0.0 c
Tomato (Castel Rock)	0.0 c	0.0 c	0.0 c
Pepper (California)	0.0 c	0.0 c	0.0 c
Bean (Giza 3)	0.0 c	0.0 c	0.0 c
Pea (Master)	0.0 c	0.0 c	0.0 c

* Figures with the same letter are not significantly different (P= 0.05).

In vitro evaluation of the inhibitory effect of chitosan and salicylic acid on the linear growth and spore germination of *F. oxysporum f.sp. niveum*:

The inhibitory effect of chitosan and salicylic acid on the linear growth and spore germination of *F. oxysporum f.sp. niveum* was evaluated under laboratory conditions. Chitosan and salicylic acid were tested at five concentrations as mentioned before. Results in Table (3) indicate that all tested concentrations of either chitosan or salicylic acid significantly reduced the linear growth and spore germination of *F. oxysporum f.sp. niveum*. Complete inhibition in linear growth was obtained with chitosan at concentration of 8.0 g/l and salicylic acid at 2.0 g/l. As for spore germination, complete inhibition was obtained with chitosan at concentration of 6.0 g/l and salicylic acid at concentration 1.5 g/l.

Effect of different concentrations of chitosan applied as seed or soil treatment on watermelon damping off and wilt diseases under greenhouse conditions:

Five concentrations of chitosan were applied either as seed or soil treatment to study their effect against Fusarium wilt incidence under greenhouse conditions.

Table 3. *In vitro* evaluation of the inhibitory effect of chitosan and salicylic acid on *F. oxysporum f.sp. niveum* linear growth and spore germination

Treatment	Concentration (g/l)	Linear growth (mm)	Reduction (%)	Spore germination (%)	Reduction (%)
Chitosan	2.0	52.5 b*	41.7	35.0 b	59.8
	4.0	27.0 c	70.0	14.0 c	83.9
	6.0	6.0 d	93.3	0.0 d	100.0
	8.0	0.0 d	100.0	0.0 d	100.0
Salicylic acid	0.5	64.0 b	28.9	42.0 b	51.7
	1.0	42.0 c	53.3	19.0 c	78.2
	1.5	14.5 d	83.9	0.0 d	100.0
	2.0	0.0 e	100.0	0.0 d	100.0
Control	0.0	90.0 a	----	87.0 a	----

* Figures with the same letter are not significantly different (P= 0.05).

Results in Table (4) show that all tested concentrations of chitosan applied either as seed or soil treatment significantly reduced the disease incidence 15 and 30 days after sowing. Soil treatment with chitosan was more efficacy than seed treatment. The highest reduction in disease incidence was obtained when the soil treated with chitosan at 6.0 and 8.0 g/kg soil which reduced the percentage of diseased plants by 60.3, 61.9 and 61.5%, respectively, for all the tested cultivars, *i.e.* cvs. Giza 1, Gorma and Aswan hybrid, respectively, when the soil was treated by 6.0 g/kg soil. Seed treatments with chitosan at concentrations of 6.0 and 8.0 g/kg soil resulted in reducing the disease incidence. Meanwhile, other concentrations were less effective

Table 4. Percentage of watermelon damping-off diseases (dead plants %) as affected by different concentrations of chitosan applied as seed or soil treatments

Chitosan (g/kg)	Dead plants (%) of watermelon cultivar								
	Giza 1			Aswan hybrid			Gorma		
	Days after sowing								
	15	30	R.%**	15	30	R.%	15	30	R.%
Seed treatment (g/l):									
2.0	18.8 b*	50.0 b	25.6	15.6 b	48.4 b	26.2	18.8 b	53.1 b	34.7
4.0	18.8 b	48.4 b	28.0	15.6 b	45.3 bc	30.9	17.2 bc	53.1 b	34.7
6.0	10.9 e	32.8 d	51.2	10.9 d	34.4 d	47.6	12.5 d	35.9 d	55.8
8.0	10.9 e	32.8 d		10.9 d	32.8 d		12.5d	35.9 d	55.8
Soil treatment (g/kg soil):									
2.0	14.1 cd	42.2 c	37.2	12.5 c	40.6 cd	38.1	17.2 bc	45.3 bc	44.3
4.0	12.5 d	37.5 c	44.2	12.5 c	35.9 d	45.3	15.6 c	40.6 c	50.1
6.0	9.4 e	26.7 de	60.3	7.8 e	25.0 e	61.9	12.5 d	31.3 e	61.5
8.0	9.4 e	23.4 e	65.2	7.8 e	25.0 e	61.9	12.5 d	28.1 e	65.4
Control 1	3.1 f	4.7 f	----	0.0 f	3.1f	----	3.1 e	3.1 f	----
Control 2	23.4 a	67.2 a	----	20.3 a	65.6 b	----	31.3 a	81.3 a	----

* Figures with the same letter are not significantly different (P= 0.05).

** Reduction (%) was recorded 30 days after sowing.

- Control 1: Non- infested soil and Control 2: Infested soil.

Effect of different concentrations of salicylic acid applied as seed or soil treatment on the incidence of watermelon wilt:

Salicylic acid solutions at various concentrations were applied either as seed or soil treatment to study their effect on Fusarium incidence of watermelon plants. Results in Table (5) indicate that all concentrations of salicylic acid applied either as seed or soil treatment significantly reduced the disease incidence. Soil treatment with salicylic acid was more efficient than seed treatments. The highest reduction was obtained when the soil was treated with salicylic acid at 1.5 and 2.0 g/kg soil which reduced the disease incidence more than 68.3, 66.6 and 70.8 % for the Giza 1, Aswan hybrid and Gorma cvs. respectively. Meanwhile, soil treatment with concentrations of 0.5 and 1.0 g/kg soil and seed treatments using 1.5 and 2.0 g/l showed moderate effect.

Table 5. Watermelon damping-off (%) as affected by different concentrations of salicylic acid applied as seed or soil treatments

Salicylic acid	Dead plants (%) of watermelon cultivar								
	Giza 1			Aswan hybrid			Gorma		
	Days after sowing								
	15	30	R.%**	15	30	R.%	15	30	R.%
Seed treatment (g/l):									
0.5	21.9 b*	39.1b	39.0	18.8 a	35.9 b	45.3	23.4 b	37.5 b	50.0
1.0	18.8 b	35.9 b	44.0	18.8 a	32.8 b	50.0	20.3 c	35.9 b	52.1
1.5	17.2 bc	28.1 c	56.2	15.5 b	26.7 c	59.3	17.2 d	29.7c	60.4
2.0	14.1 d	25.0 cd	61.0	15.5 b	26.7 c	59.3	15.6 e	29.7 c	60.4

Soil treatment (g/kg soil):									
0.5	14.1d	26.7 c	58.3	15.5 b	26.7 c	59.3	17.2 d	35.9 b	52.1
1.0	12.5 de	28.1 c	56.2	14.5 b	26.7 c	59.3	17.2 d	32.8 b	56.3
1.5	10.9 e	23.4 d	63.5	9.4 c	21.9 d	66.6	14.1 ef	25.0 c	66.7
2.0	9.4 e	20.3 d	68.3	9.4 c	21.9 d	66.6	12.5 f	21.9 c	70.8
Control 1	3.1 f	4.7 e	----	0.0 d	3.1 e	----	3.1 g	3.1 d	----
Control 2	21.9 a	64.1 a	----	20.3 a	65.6 a	----	26.7 a	75.0 a	----

* Figures with the same letter are not significantly different (P= 0.05).

** Reduction (%) was recorded 30 days after sowing.

- Control 1: Non- infested soil and Control 2: Infested soil.

Effect of soil amendment with compost for controlling Fusarium wilt of watermelon plants:

This study was carried-out in pots contained artificially infested soil with *F. oxysporum* f.sp. *niveum*. Two types, *i.e.* animal and plant composts at doses of 5.0 and 10.0 g/kg soil were tested to evaluate their effect on watermelon damping-off and/or wilt (as dead plants %). Results in Table (6) indicate that both types of compost at both the tested doses significantly reduced the disease incidence. The most effective treatment is plant compost at concentration of 10.0 g/kg soil which reduced the dead plants more than 67.4, 68.3 and 71.2% for cv. Giza 1, Aswan hybrid and cv. Gorma, respectively, 30 days after sowing. Meanwhile, other treatments showed moderate effect.

Table 6. Watermelon damping-off (%) and/or wilt diseases (as dead plants %) in response to two compost types

Compost type	Dead plants (%) of watermelon cultivar									
	Giza 1			Aswan hybrid			Gorma			
	Days after sowing									
		15	30	R.%**	15	30	R.%	15	30	R.%
Plant	5.0	10.9 c*	31.3 b	53.4	10.9 bc	25.0 b	61.0	14.1 b	28.1b	65.4
	10.0	7.8 d	21.9 c	67.4	9.4 c	20.3 c	68.3	9.4 c	23.4c	71.2
Animal	5.0	14.1 b	32.8 b	51.2	12.5 b	29.7 b	53.7	15.6 b	32.8b	59.7
	10.0	12.5 bc	28.1 b	58.2	12.5 b	25.0 b	61.0	14.1 b	28.1b	65.4
Control 1***		3.1 e	4.7 d	----	0.0 d	3.1d	----	3.1 d	3.1d	----
Control 2****		23.4 a	67.2 a	----	20.3 a	64.1a	----	31.3 a	81.3a	----

* Figures with the same letter are not significantly different (P= 0.05).

** Reduction (%) was recorded 30 days after sowing.

*** Control 1: Non- infested soil.

**** Control 2: Infested soil.

Efficacy of compost applied as soil amendment and chitosan or salicylic acid as seed bed treatments alone or in combination on Fusarium wilt incidence of watermelon transplants:

Plant compost at 10.0 g/kg soil and chitosan at 8.0 g/ kg soil or salicylic acid at 2.0 g/kg soil were tested alone or in combination to study their effect on Fusarium wilt incidence and enzyme activities of watermelon transplants.

Results in Table (7) show that plant compost, chitosan at 8.0 g/kg soil or salicylic acid at 2.0 g/kg soil applied either alone or in combination significantly reduced the disease incidence with all tested cultivars. The highest reduction was obtained by using the combined treatments between plant compost and chitosan at concentration of 8.0 g/kg soil or salicylic acid at 2.0 g/kg soil which reduced the disease incidence by 92.9, 88.5 and 86.6 % for Aswan hybrid, cvs. Gorma and Giza1, respectively. Meanwhile, single treatments showed moderate effect.

Determination of enzymes activity in watermelon transplants as affected by compost applied as soil amendment and chitosan or salicylic acid as seed bed treatments alone or in combinations:

The effect of plant compost at 10g/kg soil and chitosan at 8 g/kg soil or salicylic acid at 2 g/kg soil applied either alone or in combination on peroxidase, chitinase and β -1,3-glucanase activity was determined. Results in Table (8) reveal that all treatments increased the enzyme activities within all tested cultivars when they were applied each alone or in combination. The highest increase was obtained with combined treatments between plant compost and chitosan at concentration 8 g/kg soil or salicylic acid at 2 g/kg soil which increased the peroxidase, chitinase and β -1,3-glucanase activities more than 175.0, 200.0 and 201.0%, respectively, for all tested cultivars. Single treatments showed moderate effect. Meanwhile, plant compost was less effective.

Table 7. Fusarium wilt incidence (%) of watermelon transplants as affected by compost applied as soil amendment and chitosan or salicylic acid as seed bed treatments alone or in combinations

Treatment	Watermelon wilt incidence (%)					
	cv. Giza 1		Aswan hybrid		cv. Gorma	
	Disease incidence	R. %	Disease incidence	R. %	Disease incidence	R. %
Single treatment:						
Chitosan (8g/kg soil)	12.5 c*	73.3	12.5 c	71.5	21.9 b	60.0
SA (2g/kg soil)	12.5 c	73.3	14.1 c	67.8	21.9 b	60.0
Plant compost (10 g/kg soil)	21.9 b	53.3	21.9 b	50.0	25.0 b	54.3
Combined treatment:						
Compost + Chitosan (8g/kg soil)	6.3 d	86.6	3.1 d	92.9	6.3 de	88.5
Compost + SA (2g/kg soil)	6.3 d	86.6	3.1 d	92.9	6.3 de	88.5
Control 1**	3.1 d	----	3.1 d	----	3.1 e	----
Control 2***	46.9 a	----	43.8 a	----	54.7 a	----

* Figures with the same letter are not significantly different (P= 0.05).

** Control 1: Non- infested soil.

*** Control 2: Infested soil.

Table 8. Enzyme activities in watermelon plants as affected by compost applied as soil amendment and chitosan or salicylic acid as seed bed alone or in combination

Treatment	Increase in enzyme activities (%)								
	cv. Giza 1			Aswan hybrid			cv. Gorma		
	Po*	Ch**	β -1,3-g***	Po	Ch	β -1,3-g	Po	Ch	β -1,3-g
Single treatment									
Chitosan (8g/kg soil)	180.0	172.0	175.0	163.0	200.0	182.0	140.0	160.0	162.0
SA (2g/kg soil)	180.4	182.0	200.0	195.0	211.0	210.0	154.0	185.0	172.0
Plant compost (10 g/kg soil)	130.0	130.0	154.0	150.0	142.0	134.0	120.0	131.0	132.0
Combined treatment									
Compost + Chitosan (8g/kg soil)	192.0	222.0	214.0	194.0	210.0	220.0	175.0	200.0	201.0
Compost + SA (2 g/kg soil)	194.0	224.0	231.0	210.0	220.0	231.0	178.0	210.0	212.0
Control	5.8	1.0	2.5	6.0	1.2	2.8	4.0	0.8	1.4

* Peroxidase (Po) activity expressed as change in absorbance at 470 nm/g fresh weight/1 min.

** Chitinase (Ch) activity expressed as mM N-acetyl glucose amine equivalent released/g fresh weight/60 min.

*** β -1,3-glucanase (β -1,3-g) activity expressed as mM glucose equivalent released/g fresh weight/60 min.

Discussion

Watermelon is a widely cultivated vegetable. Fusarium wilt caused by *Fusarium oxysporum* f.sp. *niveum* is found to be a worldwide soilborne in temperate, subtropical and tropical regions (Sheng *et al.*, 2009).

In the present study, results indicated that all isolates of *F. oxysporum* f.sp. *niveum* were able to attack watermelon plants causing damping-off symptoms and wilt disease. The most aggressive isolate was Nobariya (cv. Giza 1), it caused disease incidence by 71.9, 46.9 and 51.6% for cvs. Gorma, Giza1 and Aswan hybrid, respectively, 30 days after sowing. In this respect, Nguyen (2012) reported that Fusarium wilt is one of the most severe diseases in watermelon and is caused by *F. oxysporum* f.sp. *niveum*. Among the host plants, watermelon cultivars only were susceptible to infection with the *F. oxysporum* f.sp. *niveum* (Sheng *et al.*, 2009 and Lu *et al.*, 2014).

Chitosan exhibits a variety of antimicrobial activities against plant pathogens (Benhamou, 2004; Badawy *et al.*, 2005; Abd-El-Kareem *et al.*, 2006 and El-Mohamedy *et al.*, 2013). In the present study, under laboratory experiments, results illustrated that all tested concentrations of chitosan significantly reduced the linear growth of *F. oxysporum* f.sp. *niveum*. Complete inhibition was obtained with

chitosan at concentrations of 6.0 and 8.0 g/l for linear growth and spore germination, respectively. In this respect, Kulikov *et al.* (2006) reported that the antimicrobial activity of chitosan increases with the increase in chitosan molecular weight and seems to be faster on fungi and algae than on bacteria. Fungicidal activity of chitosan has been documented against various species of fungi and Oomycetes (Vasyukova *et al.*, 2005 and Rabea *et al.*, 2005). Some of the derivatives also suppressed spore formation at rather high concentrations (Badawy *et al.*, 2005). Recently, Palma-Guerrero *et al.* (2009) demonstrated that chitosan is able to permeabilize the plasma membrane of *Neurospora crassa* and kills the cells. In general, chitosan is able to reduce the *in vitro* growth of a number of fungi and Oomycetes (Palma-Guerrero *et al.* 2008). For instance, chitosan was reported to exert an inhibitory action on the hyphal growth of numerous pathogenic fungi, including root and wilt diseases. Chitosan applied as seed or soil treatments was able to control Fusarium wilts in many plant species (Badawy *et al.*, 2005). In the present study, results indicated that all tested concentrations of chitosan significantly reduced the disease incidence. The highest reduction in disease incidence was obtained with soil treatment with chitosan at concentrations of 6.0 and 8.0 g/kg soil which reduced the disease incidence more than 60.3 and 61.9%, respectively, with all tested cultivars.

Salicylic acid is an important signalling molecule involved in both locally and systemically induced disease resistance responses. The ability to accumulate salicylic acid has been shown to be essential for systemic acquired resistance and reactions to abiotic stress in plants (Morse *et al.*, 2007; Zawoznik *et al.*, 2007; Abdel-Kader *et al.*, 2012 and Fawzy, 2013). In the present study, the inhibitory effect of salicylic acid on the linear growth and spore germination of the *F. oxysporum* f.sp. *niveum* was evaluated under laboratory conditions. All tested concentrations significantly reduced the linear growth of the fungus. Complete inhibition was obtained with salicylic acid at concentrations of 1.5 and 2.0 g/l for linear growth and spore germination, respectively. In this respect, Ozgonen *et al.*, (2001) found that salicylic acid (SA) completely inhibited the mycelial development of *F. oxysporum* f.sp. *lycopersici* at concentrations from 0.6 mM to 1.0 mM and increased dry weight of plant, length of shoot and root growth of tomato plants. Moreover, Amal (2009) reported that complete inhibition of linear growth of *Alternaria alternata*, *Fusarium oxysporum* f.sp. *phaseoli*, *F. solani* f.sp. *phaseoli*, *Macrophomina phaseolina* and *Rhizoctonia solani* was obtained with benzoic, salicylic and sorbic acids at 10 mM.

Salicylic acid is an essential component of the plant resistance to pathogens and also plays an important role in mediating plant response to some abiotic stress (Jing *et al.*, 2007). In the present study, results revealed that under greenhouse conditions, all concentrations of salicylic acid significantly reduced the disease incidence when applied as seed or soil treatments. In this respect, Suprakash and Chatterjee (2012) investigated the effect of soil application of salicylic acid (SA) and *Trichoderma harzianum* (TH) on the induction of phenolic accumulation content and defence enzymes in tomato plants infected with *F. oxysporum* f.sp. *lycopersici*. Tomato plants treated with SA showed significant decrease in wilt incidence and increase in the activities of both peroxidase and polyphenoloxidase enzymes.

Many soilborne pathogens can be reduced by application of composts made of different raw materials (Cotxarrera *et al.*, 2002) and mature composts can sustain biological control agents (Litterick *et al.* 2004 and Zhao *et al.*, 2014). In the present study, under greenhouse conditions, results indicated that two types of compost, *i.e.* animal and plant at doses of 5.0 and 10.0 g/kg soil significantly reduced the disease incidence. The most effective treatment is plant compost at 10.0 g/kg soil which reduced the disease incidence more than 67.2% for all tested cultivars. Also, plant compost at 10.0 g/kg soil and chitosan at 8.0 g/kg soil or salicylic acid at 2.0 g/kg soil were tested alone or in combination to study their effect on Fusarium wilt incidence and enzyme activities in watermelon plants. Results showed that the highest reduction was obtained with combined treatments between plant compost at 10 g/kg soil and chitosan at 8.0 g/kg soil or salicylic acid at 2.0 g/kg soil which reduced the wilt incidence more than 92.9, 88.5 and 86.6% for Aswan hybrid, cvs. Gorma and Giza1, respectively. Meanwhile, single treatments showed moderate effect. As for enzyme activities, the highest increase was obtained with combined treatments between plant compost at concentration of 10 g/kg soil and chitosan at 8.0 g / kg soil or salicylic acid at 2.0 g/ kg soil which increased the peroxidase, chitinase and β -1,3-glucanase activities more than 175.0, 200.0 and 201.0%, respectively, for all tested cultivars. Single treatments showed moderate effect.

Chitosan and salicylic acid had different properties, *i.e.* inhibitory effect against pathogenic fungus and it seems that ability to be potent elicitors of plant defence resistance. In the present study, chitosan and salicylic acid have two properties, *i.e.* antifungal activity and inducing resistance against watermelon wilt.

There are different mechanisms for reducing Fusarium wilt of watermelon plants in the present study, when chitosan was used to enhance plant defences, chitosan induced host defence responses in both monocotyledons and dicotyledons (Elwagia and Algam, 2014 and Mishra *et al.*, 2014). These responses include lignification, cytoplasmic acidification, membrane depolarization and protein phosphorylation, chitinase and glucanase activation, phytoalexin biosynthesis, generation of reactive oxygen species (Kuchitsu *et al.*, 1995), biosynthesis of jasmonic acid (Nojiri *et al.*, 1996) and the expression of unique early responsive and defence-related genes (Takai *et al.*, 2001). In addition, chitosan was reported to induce callose formation and proteinase inhibitors (Conrath *et al.*, 1989).

Moreover, it was found that SA plays its role in inducing resistance by increase the activity of chitinase, β -1,3-glucanase, peroxidase (PO), polyphenoloxidase (PPO) and phenylalanine ammonia lyase (PAL). The increase of enzymes activity was correlated with increased formation of papillae in epidermal cells (Schneider and Ullrich, 1994). Exogenous salicylic acid is able to induce antioxidant enzyme activities, formation of pathogenesis-related proteins such as β -1,3-glucanase and chitinase, and expression of antioxidant enzyme genes in some plant leaves (Fernandes *et al.*, 2006 and Chen *et al.*, 2006). Furthermore, as for compost treatments Hoitink and Boehm (1999) have postulated the following biological mechanisms of disease control with composts: parasitism against pathogens by beneficial microorganisms, antibiotic production by beneficial microorganisms, competition for nutrients by beneficial microorganisms, activation of disease-

resistance genes in plants by microorganisms and improved plant nutrition and vigour, leading to enhance disease resistance. The latter two modes of action have been used to explain instances where disease control resulting from compost amendment of soil was not accompanied by a corresponding reduction in pathogen inoculum. Also, several studies under controlled conditions have demonstrated a suppressive effect of composts on soilborne diseases such as damping-off, root rot, and wilt (Hoitink and Boehm, 1999).

References

- Abdel-Kader, M.M.; El-Mougy, Nehal S.; Abdalla, M.A.; Abd-El-Kareem, F. and Fatooh, Y.O. 2012. Effect of some plant resistance inducers, essential oils and plant extracts on antagonistic ability of bacteria and yeast bio-agents *in vitro*. *J. Appl. Sci. Res.*, **8**(1): 543-555.
- Abd-El-Kareem, F.; El-Mougy, Nehal S.; El-Gamal, Nadia G. and Fatooh, Y.O. 2006. Use of chitin and chitosan against tomato root rot disease under greenhouse conditions. *Res. J. Agric. and Biol. Sci.*, **2**(4): 164-169.
- Abd-El-Kareem, F. and Abd-El-Latif, Faten M. 2012. Using bicarbonates for controlling late blight disease of potato plants under field conditions. *Life Sci. J.*, **9**(4): 2080-2085
- Abd-El-Kareem, F.; Abdel-Kader, M.M.; Fatooh, Y.O.; Abd-Alla, M.A. El-Mougy, Nehal S.; El-Mohamedy, R.S. and El-Gamal, N.G. 2013a. Induction of systemic resistance in faba bean against chocolate spot diseases severity using chemical inducers under field conditions. *J. Appl. Sci. Res.*, **9**(6): 4006-4014.
- Abd-El-Kareem, F.; El-Mougy, Nehal S. and Abdel-Kader, M.M. 2013b. Application of compost and bio-agents as integrated soil treatment for controlling peanut crown rot disease under field conditions. *Advances in Agric. Sci. and Engineer. Res.*, **3**(5): 858-866.
- Abeles, F.B. and Forrence, L.E. 1970. Temporal and hormonal control of β -1,3-glucanase in *Phaseolus vulgaris* L. *Plant Physiol.*, **45**: 395-400.
- Abeles, F.B.; Bosshart, R.P.; Forrence, L.E. and Habig, W.H. 1971. Preparation and purification of glucanase and chitinase from bean leaves. *Plant Physiol.*, **47**: 129-134.
- Amal, A.H.H. 2009. Control of bean root-rot diseases by using resistance chemical inducers. *Alex. J. Agric. Res.*, **54**: 175-187.
- Badawy, M.T.; Rogge, T.M.; Stevens, C.V.; Höfte, M.; Steurbaut, W. and Smagghe, G. 2005. Insecticidal and fungicidal activity of new synthesized chitosan derivatives. *Pest Management Sci.*, **61**: 951-960.
- Benhamou, N. 2004. Potential of the mycoparasite, *Verticillium lecanii*, to protect citrus fruit against *Penicillium digitatum*, the causal agent of green mould: A comparison with the effect of chitosan. *Phytopathology*, **94**:693-705.

- Booth, C. 1971. *The Genus Fusarium*. Commonweals Mycol. Inst., Kew, Surrey, England, 235pp.
- Chen, J.Y.; Wen, P.F.; Kong, W.F.; Pan, Q.H.; Zhan, J.C. and Huang, W.D. 2006. Effect of salicylic acid on phenylpropanoids and phenylalanine ammonialyase in harvested grape berries. *Posthar. Biol. Technol.*, **40**: 64-72.
- Conrath, U.; Domard, A. and Kauss, H. 1989. Chitosan-elicited synthesis of callose and of coumarin derivatives in parsley cell suspension cultures. *Plant Cell Rep.*, **8**: 152-155.
- Cotxarrera, L.; Trillas-Gay, M.I.; Steinberg, C. and Alabouvette, C. 2002. Use of sewage sludge compost and *Trichoderma asperellum* isolates to suppress Fusarium wilt of tomato. *Soil Biol. Biochem.*, **34**: 467-476.
- Elad, Y. and Baker, R. 1985. Influence of trace amount of cations and siderophore-producing *Pseudomonads* on chlamydospores of *Fusarium oxysporum*. *Phytopathology*, **75**: 1047-1052.
- El-Mohamedy, R.S.; Abdel-Kader, M.M.; Abd-El-Kareem, F. and El-Mougy, Nehal S. 2013. Inhibitory effect of antagonistic bio-agents and chitosan on the growth of tomato root rot pathogens *in vitro*. *J. Agric. Technol.*, **9**:1521-1533.
- Elwagia, M.E.F. and Algam, S. 2014. Evaluation of chitosan efficacy on tomato growth and control of early blight disease. 3rd Conf. Pests Manage. in Sudan, CPRC-ARC, Wad Median (Sudan), p. 10.
- Fawzy, A.M. 2013. Improvement of biocontrol of damping-off and root rot / wilt of faba bean by salicylic acid and hydrogen peroxide. *Mycobiology*; **41**: 47-55.
- Fernandes, C.F.; Silveira, J.A.; Vasconcelos, G.; Oliveira, I.M. and Abreu, J.T. 2006. Induction of an anionic peroxidase in cowpea leaves by exogenous salicylic acid. *J. Plant Physiol.*, **163**: 1040-1048.
- Gilman, J.C. 1957. *A Manual of Soil Fungi*. The Iowa State College Press Ames. Iowa. 450 pp .
- Goldschmidt, E.E.; Goren, R. and Monselise, S.P. 1968. The IAA oxidase system of citrus roots. *Planta*, **72**: 213-222.
- Hoitink, H.A.J. and Boehm, M.J. 1999. Biocontrol within the context of soil microbial communities: A substrate-dependent phenomenon. *Annu. Rev. Phytopathol.*, **37**: 427-446.
- Jing, C.; Zhu, C.; Li, L.P.; Sun, Z.Y. and Pan, X.B. 2007. Effects of exogenous salicylic acid on growth and H₂O₂-metabolizing enzymes in rice seedlings under lead stress. *J. Environ. Sci.*, **19**: 44-49.
- Kuchitsu, K.; Kosaka, H.; Shiga, T. and Shibuya, N. 1995. Evidence for generation of hydroxyl radical triggered by *N* – acetyl chitoooligo saccharide elicitor and a protein phosphatase inhibitor in suspension-cultured rice cells. *Protoplasma*, **88**: 138-142.

- Kulikov, S.N.; Chirkov, S.N.; Lopatin, S. and Varlamov V.P. 2006. Effect of the molecular weight of chitosan on its antiviral activity in plants. *Prik Biokhim Mikrobiol.*, **42**(2): 224-228.
- Litterick, A.; Harrier, L.; Wallace, P.; Watson, C. and Wood, M. 2004. The role of uncomposted materials, composts, manures and compost extracts in reducing pest and disease incidence and severity in sustainable temperate agricultural and horticultural crop production. *Rev. Plant Sci.*, **23**: 453-479
- Lu, G.; Guo, S.; Zhang, H.; Geng, L.; Martyn, M.D. and Xu, Y. 2014. Colonization of Fusarium wilt-resistant and susceptible watermelon roots by a green-fluorescent – protein – tagged isolate of *Fusarium oxysporum* f.sp. *niveum*. *J. Phytopathol.*, **2**: 228-237.
- Mishra, S.; Jagadeesh, K.S.; Krishnaraj, P.U. and Prem, S. 2014. Biocontrol of tomato leaf curl virus in tomato with chitosan supplemented formulations of *Pseudomonas* sp. under field conditions. *Austral. J. Crop Sci.*, **8**: 347-355.
- Monreal, J. and Reese, E.T. 1969. The chitinase of *Serratia marcescens*. *Canad. J. Microbiol.*, **15**: 689-696.
- Morse, A.M.; Tschaplinski, T.J.; Dervinis, C.; Pijut, P.M.; Schmelz, E.A.; Day, W. and Davis, J.M. 2007. Salicylate and catechol levels are maintained in nahG transgenic poplar. *Phytochemistry*, **68**: 2043-2052.
- Neler, J.; Wassermann, W. and Kutner, M.H. 1985. Applied linear statistical models. Pages: 117-155. In: *Regression Analysis of Variance and Experimental Design*. 2nd Ed. D. Richard (ed.). Irwin Inc., Homewood Illinois, USA.
- Nguyen, L.T. 2012. Outbreak of Fusarium wilt in seedless watermelon seedlings in the Northern Territory, Australia. *Austral. Plant Dis. Notes*. doi: 10.1007/s13314-012-0053.
- Nojiri, H.; Sugimori, M.; Yamane, H.; Nishimura, Y. and Ohmori, T. 1996. Involvement of jasmonic acid in elicitor-induced phytoalexin production in suspension-cultured rice cells. *Plant Physiol.*, **110**: 387-392.
- Ozgonen, H.; Bicici, M. and Erklc, A. 2001. The effect of salicylic acid and endomycorrhizal fungus *Glomus etunicatum* on plant development of tomatoes and Fusarium wilt caused by *Fusarium oxysporum* f.sp. *lycopersici*. *Turkish J. Agric. and Forestry*, **25**: 25-29.
- Palma-Guerrero, J.; Huang, I.C.; Jansson, H.B.; Salinas, J.; Lopez, L.V. and Read, N.D. 2008. Effect of chitosan on hyphal growth and spore germination of plant pathogenic and biocontrol fungi. *J. Appl. Microbiol.*, **104**: 541-553.
- Palma-Guerrero, J.; Huang, I.C.; Jansson, H.B.; Salinas, J.; Lopez, L.V. and Read, N.D. 2009. Chitosan permeabilizes the plasma membrane and kills cells of *Neurospora crassa* in an energy dependent manner. *Fungal Gen. Biol.*, **46**: 585-594.

- Rabea, M.T.; Rogge, T.M.; Stevens, C.V. and Smagghe, G. 2005. Insecticidal and fungicidal activity of new synthesized chitosan derivatives. *Pest Management Sci.*, **61**: 951-960
- Ried, J.D. and Ogryd-Ziak, D.M. 1981. Chitinase over producing mutant of *Servatia marcescens*. *Appl. and Environ. Microbiol.*, **41**: 664-669.
- Schneider, S. and Ullrich, W.R. 1994. Differential induction of resistance and enhanced enzyme activities in cucumber and tobacco caused by treatment with various abiotic and biotic inducers. *Physiol. and Mole. Plant Pathol.*, **45**: 291-304.
- Sheng, H.; Ning, X.; Qin, J. and Miao, W. 2009. Suppression of Fusarium wilt of watermelon by a bio-organic fertilizer containing combinations of antagonistic microorganisms. *Biocontrol*, **54**: 287-300.
- Suprakash, O. and Chatterjee, N.C. 2012. Induction of resistance in tomato plants against *Fusarium oxysporum* f.sp. *lycopersici* mediated through salicylic acid and *Trichoderma harzianum*. *J. Plant Protect. Res.*, **52**: 220-225
- Takai, R.; Hasegawa, K.; Kaku, K.; Shibuya, N. and Minami, E. 2001. Isolation and analysis of expression mechanisms of a rice gene, *EL5*, which shows structural similarity to *ATL* family from *Arabidopsis*, in response to *N*-acetylchitooligosaccharide elicitor. *Plant Sci.*, **160**: 577-583.
- Trillas, M.I.; Casanova, E.; Corxarrera, L.; Ordovas, J.; Borrero, C. and Aviles, M. 2006. Composts from agricultural waste and the *Trichoderma asperellum* strain T-34 suppress *Rhizoctonia solani* in cucumber seedlings. *Biol. Control*, **39**: 32-38.
- Tziros, G.T.; Lagopodi, A. L. and Tzavella-Klonari, K. 2007. Reduction of Fusarium wilt in watermelon by *Pseudomonas chlororaphis* PCL 1391 and *P. fluorescens* WCS365. *Phytopathol.Mediterr.* **46**, 320 – 323.
- Vasyukova, N.I.; Chalenko, G.I.; Gerasimova, N.G.; Perekhod, E.A.; Ozeretskovskaya, O.L. and Albulov, A.I. 2005. Chitin and chitosan derivatives as elicitors of potato resistance to late blight. *Appl. Biochem. Microbiol.*, **36**: 372-376.
- Zawoznik, M.S.; Benavides, M.P. and Tomaro, M.L. 2007. Endogenous salicylic acid potentiates cadmium-induced oxidative stress in *Arabidopsis thaliana*. *Plant Sci.*, **173**: 190-197.
- Zhao, S.; Liu, D.; Ling, N.; Chen, F.; Fang, W. and Shen, Q. 2014. Bio-organic fertilizer application significantly reduced the *Fusarium oxysporum* population and alters the communities of watermelon Fusarium wilt rhizosphere soil. *Biol. Fertil. Soil, On line*, DOI:10.1007/00374-014.

(Received 12/06/2014;
in revised form 15/07/2014)

المقاومة المستحثة في نباتات البطيخ ضد مرض الذبول الفيوزاريومي باستخدام المستحضات الكيماوية والكمبوست تحت ظروف الصوبة

عبد مهيدي محمد مهيدي* ، محمد هارون عبد المجيد* ،
فاتن محمود عبد اللطيف* ، محمد محمد دياب** ،
نهال محمد سيد**

* قسم أمراض النبات - كلية الزراعة - جامعة بنها- مصر.
** قسم أمراض النبات - المركز القومي للبحوث - مصر.

أحدثت كل العزلات المختبرة من الفطر *Fusarium oxysporum* أعراض الذبول وموت البادرات (سقوط البادرات) علي نباتات البطيخ وكانت اقوي عزلة هي النوبارية (معزولة من صنف البطيخ جيزة 1) حيث أدت الي نسبة اصابة بمقدار 92.09 و 88.5 و 86.6% بالنسبة لاصناف البطيخ هجين أسوان وجورما و جيزة 1 علي الترتيب.

لم تسبب عزلة الفطر *F. oxysporum* أي اصابة علي أي أنواع أو أصناف نباتية أخرى من الفصيلة القرعية او الفصيلة الباذنجانية أو البقولية الا أصناف نباتات البطيخ فقط مما يدل علي أنها عزلة من الفطر *F. oxysporum* f.sp. *niveum*. أدت كل تركيزات الكيتوزان وحمض الساليسيلك و الكمبوست النباتي والحيواني الي انخفاض نسبة حدوث مرض الذبول الفيوزاريومي في نباتات البطيخ .

تم دراسة تكامل المعاملات بين معاملة التربة في الاصلص المعداه بالفطر الممرض بالكمبوست النباتي بتركيز 10 جم/كجم تربة و معاملة مهد البذرة لشتلات البطيخ بالكيتوزان بتركيز 8 جم/كجم تربة وحمض الساليسيلك بتركيز 2 جم/كجم تربة علي نسبة حدوث مرض الذبول الفيوزاريومي علي نباتات البطيخ وكذلك نشاط الانزيمات المسئولة عن المقاومة في نباتات البطيخ وأوضحت النتائج ما يلي:-

أدي تكامل المعاملات بين معاملة التربة بالكمبوست النباتي بتركيز 10 جم/كجم تربة و معاملة مهد البذرة لشتلات البطيخ بالكيتوزان بتركيز 8 جم / كجم تربة وحمض الساليسيلك بتركيز 2 جم/كجم تربة الي مكافحة فعالة لمرض الذبول وموت البادرات علي نباتات البطيخ حيث أدت الي انخفاض نسبة حدوث المرض بأكثر من 86.3 و 92.8 و 88.3% بالنسبة لاصناف البطيخ جورما و جيزة 1 و هجين أسوان علي الترتيب. بالنسبة لنشاط الانزيمات أدت المعاملات السابق ذكرها الي زيادة معنوية في كل الانزيمات المختبرة حيث أدت الي زيادة مقدارها 175.0 و 200.0 و 210.0% لانزيمات البيروكسيداز والكيتيناز والبيتا جلوكانيز علي الترتيب مع كل الاصناف المختبرة.